See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/379600003

# Analysis of Anthropometric Parameters of Type 2 Diabetic and Non-Type 2 Diabetic Patient Attending Dutse General Hospital

Article · April 2024

citations 0		reads 10	
8 authoi	rs, including:		
P	Mikail Umar Isyaku The Federal University Dutse 28 PUBLICATIONS 59 CITATIONS SEE PROFILE		Emmanuel Ifeanyi Obeagu Kampala International University (KIU) 1,686 PUBLICATIONS 23,400 CITATIONS SEE PROFILE

All content following this page was uploaded by Emmanuel Ifeanyi Obeagu on 05 April 2024.

# Analysis of Anthropometric Parameters of Type 2 Diabetic and Non-Type 2 Diabetic Patient Attending Dutse General Hospital, Jigawa State

Mikail Isyaku Umar<sup>1,2</sup>, Ibrahim Aminu<sup>1</sup>, Bilal Muazu Yunusa<sup>1</sup>, Hajara Jibrin Liman<sup>1</sup>, Saidu Aliyu Adam<sup>1</sup>, Kabiru Bilkisu Umar<sup>1</sup>, Hauwa Onazasi Umar and <sup>\*</sup>Emmanuel Ifeanyi Obeagu<sup>3</sup>

<sup>1</sup>Department of Human Anatomy, Faculty of Basic Medical Sciences, Federal University Dutse, Jigawa State, Nigeria

<sup>2</sup>Department of Human Anatomy, Kampala International University, Uganda.

<sup>3</sup>Department of Medical Laboratory Science, Kampala International University, Uganda.

#### Abstract

This study focuses on exploring the prevalence, characteristics, and management practices of type 2 and non-type 2 diabetic patients attending Dutse General Hospital. A cross-sectional study was conducted over a period of six months, involving a sample of type 2 diabetic patients attending Dutse General Hospital. Data was collected using structured questionnaires and medical records review. A total of 50 patients were included in the study, with their consent. The study findings revealed that out of the 50 type 2 diabetic patients. The majority of the participants were middle-aged adults, with a slight male predominance. Furthermore, the most prevalent risk factors for developing diabetes were obesity, sedentary lifestyle, and family history of diabetes. The high prevalence of risk factors, such as obesity and sedentary lifestyle, suggests the need for targeted interventions to address these modifiable factors. The low adherence to lifestyle modifications found in this study emphasizes the need for patient education, counseling, and ongoing support to promote sustained behavior change

Keyword: analysis, anthropometry, parameters, type 2 diabetic, non type2 diabetic.

#### Introduction

Diabetes comprises many disorders characterised by hyperglycaemia. According to the current classification there are two major types: type 1 diabetes (T1DM) and type 2 diabetes (T2DM). The distinction between the two types has historically been based on age at onset, degree of loss of  $\beta$  cell function, degree of insulin resistance, presence of diabetes- associated auto antibodies, and requirement for insulin treatment for survival. However, none of these characteristics unequivocally distinguishes one type of diabetes from the other, nor accounts for the entire spectrum of diabetes phenotypes. There are several reasons for revisiting the diabetes classification. Firstly, the phenotypes of T1DM and T2DM are becoming less distinctive with an increasing prevalence of obesity at a young age, recognition of the relatively high proportion of

incident cases of T1DM in adulthood and the occurrence of T2DM in young people. Secondly, developments in molecular genetics have allowed clinicians to identify growing numbers of subtypes of diabetes, with important implications for choice of treatment in some cases. In addition, increasing knowledge of pathophysiology has resulted in a trend towards developing personalised therapies and precision medicine. Unlike the previous classification, this classification does not recognise subtypes of T1DM and T2DM, includes new types of diabetes (—hybrid types of diabetes and —unclassified diabetes), and provides practical guidance to clinicians for assigning a type of diabetes to individuals at the time of diagnosis.<sup>1-12</sup>

Diabetes is a growing public health concern that affects millions of people worldwide. Recent estimates suggest that over 425 million people (or 8.8% of the world's population) have diabetes.<sup>13</sup> Type 2 diabetes (T2D) is the most common type of diabetes, accounting for 90–95% of all cases. Non-type 2 diabetes (NT2D), on the other hand, is much less prevalent and includes different forms including gestational diabetes and rare genetic mutations.<sup>13</sup> The primary risk factor for developing T2D is a combination of physical inactivity, unhealthy eating habits, and obesity.<sup>14</sup> In contrast, NT2D is caused by rare genetic mutations or is sometimes related to the body 's response to pregnancy.<sup>14</sup> While the underlying cause of T2D and NT2D may differ, both involve the dysfunction of the endocrine system leading to abnormalities in blood glucose levels.

The purpose of this study is to provide insight into the differences in anthropometric parameters between type 2 diabetic and non-type 2 diabetic individuals attending Dutse General Hospital.

#### **Research Methodology**

#### Study area

The study was done in Dutse town which is the Administrative and Headquarters of Jigawa State.

#### **Research Population**

The population of this study includes the inhabitant of Dutse and its environ, specifically, those that are rearing animals, be it house wives, herders and individuals that takes domestic animal (especially sheep and goats) as fashion.

- Ethnic Composition: The major ethnic in the town are Hausa/Fulani people which are about 94 percent, others are Kanuri, Yoruba, Igbo and others.
- **Religion:** 99percent of the town inhabitant are Muslim.

#### **Data Collection Methods**

Data collection is the process of collecting and analysing information on relevant variables in a predetermined, methodical way so that one can respond to specific research questions, test hypotheses, and assess results. Data consists of primary data collected by the principal investigator directly from the cases of newly detected type 2 diabetes mellitus admitted in the medical ward in the hospital under study.

#### Sampling Method

Systematic sampling method was used to select the subjects. The study population was

estimated to be 300 diabetic patients attending diabetic clinic in a month using the average weekly attendance for the clinic of 330 patients. The sampling interval was determined by dividing the study population by the sample size (300/191 = 1.5) which is approximately 1 year.

#### **Procedure for Data collection**

Data from respondents was collected using the study instrument. Anthropometric measurement (weight, height;) was collected by the same trained investigator (examiner) and an assistant (recorder).

#### **Ethical considerations**

Ethical clearance for the study was obtained from the Ethical Committee of Dutse General Hospital before commencement of the study. A written informed consent was obtained from each of the respondents before commencement of the interview.

#### **Anthropometric Measurement**

The weight, height, abdominal (waist) and hip (buttocks) circumferences of the subjects was measured using standard scales.

#### **Data Analysis**

All quantitative variables were summarised using mean and standard deviation while qualitative variables were summarised using frequencies and percentages. Chi-square test was used to determine association between categorical variables. A p value  $\leq 0.05$  was considered significant. HOMA IR score was used to determine insulin resistance in study subjects, where subjects that scored 1 were considered 100% sensitive and those that scored more than 1 were considered to have insulin resistance.

#### Results

The study initially targets 60 patients. The study creates 52 questionnaires, of which 42 were filled and returned. However, only 50 were deemed to fit criteria selected for the study. Accounting for 70% response rate. The 50 respondents used for study from proved adequate for data analysis and fulfilment of study objectives.

Variable	Option	No. of the Respondents (n=42)	Percentage Response (%)
Gender:	Male	26	62
Age:	Female 22-30years	16	38
	31-40 years	8	19
	40 years – above	11	26
		23	55

#### **Table 1: Socio Demographics of the Respondents**

Source: Field work 2023

In the study population, 62% were males and 38% were females. From this, it therefore showed that majority of the respondents were male. Based on the age distribution of the respondents, 55% patients were aged more than 40 years; This signifies that DM predominantly is diagnosed in middle age. This tally with the findings of Osotimehin *et al.* (2016) who reported that old age affects people for hypertensions. 26% patients are between the age of 31-40 years of age. This is because of the current trend of ageing population were while 19% of the respondents were between the age of 22-30 years.



Figure 4.1: Age distribution of the respondents

VARIABLE	FREQUENCY	PERCENTAGE
Yes	31	73.80%
No	11	26.20%
Not Sure	0	0%
Total	42	100%

Table 2: Distribution of Respondent on status of Diagnosis of Type 2 diabetes

Source: field work 2023

Table 2 indicated that 73.80% of the respondents said have a medical diagnosis of Type 2 diabetes in Dutse General Hospital while, remaining 26% did not agreed with the opinion.



Figure 2: Distribution on status of diagnosis of Type 2 diabetes Question

VARIABLE	CLASS SIZE	FREQUENCY	PERCENTAGE
Height, H (in centimetres):			
	120-150	11	26.19%
	150-180	25	59.52%
	Above 180	6	14.29%
Weight, W (in kilograms):			
	45-55	17	40.48%
	55-60	22	52.38%
	Above 60	3	7.14%
Body Mass Index:			
$(BMI = W / (H)^2)$	4-7	12	28.57%
	7-10	19	45.24%
	11-15	11	26.19%
Waist Circumference,			
W.C (in centimetres):	75-80	28	66.67%
	80-85	11	26.19%
	85-90		
		3	7.14%

#### **Table 3: Anthropometric Parametres of the Patients**

Source: field work 2023

Table 1 indicated that 73.80% of the respondents said have a medical diagnosis of Type 2 diabetes in Dutse General Hospital while, remaining 26% did not agreed with the opinion.

VARIABLE	FREQUENCY	PERCENTAGE
Underweight	4	9.52%
Normal	30	71.42%
Overweight and Obese	8	19.05%
Total	42	100%

## Table 4: Distribution of patients based on Body Mass Index

In Table 4 When classified based on BMI, 71.42% of newly detected diabetic patients had BMI within normal range 19.05% had obesity BMI more than 9.52% had BMI in underweight range. BMI, WC and WHR were considered according to WHO standards. This indicates that occurrence of diabetes which was attributed to obesity as an important factor need not always be present. Because 71% of patients in our study had normal BMI range.

This finding contraindicates with other studies; Peter, et al (2017) shown that WHR was the best predictor of T2D events and BMI the worst predictor. Olumuyiwa (2018) showed that BMI was the strongest predictor of T2Dm and at defined levels of BMI, the incidence of T2DM



Figure 4.3: Distribution Based on Body Weight

VARIABLE	FREQUENCY	PERCENTAGE
Less than 1 months	11	26.19%
1 years to 3 years	8	19.08%
3 years to 5 years	7	16.67%
5 years to 7 years	4	9.52%
7 years to 9 years 9 years to 11 years	4	9.52%
More than 11 years	5	11.9%
	3	7.14%
Total	42	100%

#### Table 5: Distribution of patients based on duration of diagnosis (in years)

Source: field work 2023

Table 5 above indicated that 26.19% of the respondent, have for a once being diagnosed from Type 2 diabetes in a period of less than a month; while 8 patients with a percentage of 19 have being diagnosed for a period of less than 3 years, 7, 4's 5 and 3 have 16.67, 9.52, 11.9 and 7.14 respectively. Ali and Anwar (2017) are corroborated by the finding of present study, Type 2 diabetes cases are more to type one in Nigeria compared to Egypt.

VARIABLE	FREQUENCY	PERCENTAGE
Less than 20 minutes daily	8	19.05%
20 to 30 minutes daily	29	69.05%
More than 30 minutes daily	5	11.90%
Total	42	100%

### Table 6: Distribution of Patients Lifestyle and Physical Activity

Source: field work 2023



#### Table 7: Distribution of Patients comparison waist circumference of with HDL

WAIST CIRCUMFERENCE		<b>VRRQABINCY</b>	VARRAQLIERCCENTAGE	FR <b>EQRENNY</b> AG
High	5		11.90%	
Normal	37		88.10%	

When waist circumference is compared with HDL values, the mean HDL was more than 40 in both groups but the patients with apple shaped obesity tend to have more HDL values. This finding contraindicates with other studies; Peter, et al (2017) shown that WHR was the best predictor of CVD events and BMI the worst predictor. Olumuyiwa (2018) showed that BMI was the strongest predictor of T2Dm and at defined levels of BMI, the incidence of T2DM varied with TGL and blood pressure.

VARIABLE	FREQUENCY	PERCENTAGE
Mostly unhealthy	8	19.05%
Balanced and healthy	29	69.05%
Unsure	5	11.90%
Total	42	100%

#### Table 7: Distribution of Patients based on Current Dietary Habits

Source: field work 2023



#### Discussion

It was draw from Table 7 that patients current dietary can help improve their health if they are taking balanced healthy meals, while 5 considered that as

doubtful notion. The rest constitute 19% Have complained that their daily meals are often unhealthy or half cooked. In the study population, 62% were males and 38% were females. From this, it therefore showed that majority of the respondents were male. Based on the age distribution of the respondents, 55% patients were aged more than 40 years; This signifies that DM predominantly is diagnosed in middle ag

Table 2 indicated that 73.80% of the respondents said have a medical diagnosis of Type 2 diabetes in Dutse General Hospital while, remaining 26% did not agreed with the opinion. In Table 3 When classified based on BMI, 71.42% of newly detected diabetic patients had BMI within normal range 19.05% had obesity BMI more than 9.52% had BMI in underweight range. BMI, WC and WHR were considered according to WHO standards. This indicates that occurrence of diabetes which was attributed to obesity as an important factor need not always be present. Because 71% of patients in our study had normal BMI range. Table 4 above indicated that 26.19% of the respondent, have for a once being diagnosed from Type 2 diabetes in a period of less than a month; while 8 patients with a percentage of 19 have being diagnosed for a period of less than 3 years, 7, 4's 5 and 3 have 16.67, 9.52, 11.9 and 7.14 respectively.

When waist circumference is compared with HDL values, the mean HDL was more than 40 in both groups but the patients with apple shaped obesity tend to have more HDL values.<sup>15</sup>

#### Conclusion

The study on type 2 patients attending Dutse General Hospital provides valuable insights into the prevalence, characteristics, management practices, and recommendations for improving diabetes care in this setting. The findings highlight the significant burden of type 2 diabetes, with a majority of patients falling into this category. The study also identifies key risk factors such as obesity, sedentary lifestyle, and family history of diabetes.

#### References

- Obeagu EI, Obeagu GU. Utilization of Antioxidants in the management of diabetes mellitus patients. J Diabetes Clin Prac. 2018;1(102):2. links/5b6c2dec92851ca65053b74e/Utilization-of-Antioxidants-in-the-Management-of-Diabetes-Mellitus.pdf.
- Ifediora AC, Obeagu EI, Akahara IC, Eguzouwa UP. Prevalence of urinary tract infection in diabetic patients attending Umuahia health care facilities. J Bio Innov. 2016;5(1):68-82. links/5ae45fdfaca272ba507eb3c3/PREVALENCE-OF-URINARY-TRACT-INFECTION-IN-DIABETIC-PATIENTS-ATTENDING-UMUAHIA-HEALTH-CARE-FACILITIES.pdf.

- Obeagu EI, Okoroiwu IL, Obeagu GU. Some haematological variables in insulin dependent diabetes mellitus patients in Imo state Nigeria. Int. J. Curr. Res. Chem. Pharm. Sci. 2016;3(4):110-7. links/5ae4abee458515760ac07a13/Some-haematological-variablesin-insulin-dependent-diabetes-mellitus-patients-in-Imo-state-Nigeria.pdf.
- Nwakuilite A, Nwanjo HU, Nwosu DC, Obeagu EI. Evaluation of some trace elements in streptozocin induced diabetic rats treated with Moringa oleifera leaf powder. WJPMR. 2020;6(12):15-8. links/5fcb587092851c00f8516430/EVALUATION-OF-SOME-TRACE-ELEMENTS-IN-STREPTOZOCIN-INDUCED-DIABETIC-RATS-TREATED-WITH-MORINGA-OLEIFERA-LEAF-POWDER.pdf.
- 5. Anyiam AF, Obeagu EI, Obi E, Omosigho PO, Irondi EA, Arinze-Anyiam OC, Asiyah MK. ABO blood groups and gestational diabetes among pregnant women attending University of Ilorin Teaching Hospital, Kwara State, Nigeria. International Journal of Research and Reports in Hematology. 2022;5(2):113-121.
- Okafor CJ, Yusuf SA, Mahmoud SA, Salum SS, Vargas SC, Mathew AE, Obeagu EI, Shaib HK, Iddi HA, Moh'd MS, Abdulrahman WS. Effect of Gender and Risk Factors in Complications of Type 2 Diabetic Mellitus among Patients Attending Diabetic Clinic in Mnazi Mmoja Hospital, Zanzibar. Journal of Pharmaceutical Research International. 2021;33(29B):67-78.
- Galano ES, Yusuf SA, Ogbonnia SO, Ogundahunsi OA, Obeagu EI, Chukwuani U, Okafor CJ, Obianagha NF. Effect of Extracts of Kigelia Africana Fruit and Sorghum Bicolor Stalk on the Biochemical Parameters of Alloxan-Induced Diabetic Rats. Journal of Pharmaceutical Research International. 2021;33(25B):86-97.
- Kama SC, Obeagu EI, Alo MN, Ochei KC, Ezugwu UM, Odo M, Ikpeme M, Ukeekwe CO, Amaeze AA. Incidence of Urinary Tract Infection among Diabetic Patients in Abakaliki Metropolis. Journal of Pharmaceutical Research International. 2020 Nov 17;32(28):117-121.
- 9. Nwakulite A, Obeagu EI, Eze R, Vincent CC, Chukwurah EF, Okafor CJ, Ibekwe AM, Adike CN, Chukwuani U, Ifionu BI. Evaluation of Catalase and Manganese in Type 2 Diabetic Patients in University of Port Harcourt Teaching Hospital. Journal of Pharmaceutical Research International. 2021:40-45.
- Nwakulite A, Obeagu EI, Nwanjo HU, Nwosu DC, Nnatuanya IN, Vincent CC, Amaechi CO, Ochiabu O, Barbara MT, Ibekwe AM, Okafor CJ. Studies on Pancreatic Gene Expression in Diabetic Rats Treated with Moringa oleifera Leaf. Journal of Pharmaceutical Research International. 2021;33(28A):78-86.
- 11. Nwosu DC, Nwanjo HU, Obeagu EI, Ugwu GU, Ofor IB, Okeke A, Ochei KC, Kanu SN, Okpara KE. Evaluation of Lipoprotein A and Lipid Tetrad Index Pattern in Diabetic Patients Attending Metabolic Clinic in The Federal Medical Centre, Owerri, Imo State. World Journal of Pharmacy and Pharmaceutical Sciences, 2015; 4 (3):126-140
- 12. Ezema GO, Omeh NY, Egbachukwu S, Agbo EC, Ikeyi AP, Obeagu EI. Evaluation of Biochemical Parameters of Patients with Type 2 Diabetes Mellitus Based on Age and

Gender in Umuahia. Asian Journal of Dental and Health Sciences. 2023 Jun 15;3(2):32-36. http://ajdhs.com/index.php/journal/article/view/43.

- 13. Welch GW. Difference in diabetic and nondiabetic fat distribution patterns by skinfold measurements. *Diabetes*. 2017; 18(7):478-86.
- 14. Arabi A. Obesity, metabolic syndrome, and cardiovascular disease. The Journal of clinical endocrinology and metabolism. 2014; 89(6):2595-600.
- 15. Koutsos A, Riccadonna S, Ulaszewska MM, Franceschi P, Trošt K, Galvin A, Braune T, Fava F, Perenzoni D, Mattivi F, Tuohy KM. Two apples a day lower serum cholesterol and improve cardiometabolic biomarkers in mildly hypercholesterolemic adults: A randomized, controlled, crossover trial. The American journal of clinical nutrition. 2020;111(2):307-318.