















SYSTEMATIC REVIEW

Plant-based management of Iron Deficiency Anemia (IDA) in East Africa community: a systematic review

[version 1; peer review: awaiting peer review]

Robert Siida ¹, Ilemobayo Victor Fasogbon ¹, Angela Mumbua Musyoka ¹, Basil Uchechukwu Nwali², Sandra Omo Etumah¹, Idara Asuquo Okon³, Ejike Daniel Eze³, Afodun Adam⁴, Herbert Mbyemeire¹, Comfort Danchal Vandu⁵, Elna Owembabazi¹, Michael Ben Okon ¹, Peter Chinedu Agu ^{6,7}, Usman Michael Ibe ⁸, Wusa Makena⁸, Augustine Oviosun⁸, Monday Ekon Etukudo ⁸, Regan Mujinya³, Reuben Samson Dangana ¹, Daniel Ejim Uti ⁹, Vivian Onyinye Ojiakor⁸, Esther Ugo Alum ⁹, Swase Dominic Terkimbi ¹, Lucy Aja ¹⁰, Sanusi Ahmed Jega¹, Ugwu Okecukwu Paul Chima⁹, Nancy Bonarare Mitaki ¹, Josiah Eseoghene Ifie ¹, Mukangendo Mecthilde⁴, Sinbad Olubukola Olorunnisola¹, Patrick Maduabuchi Aja ^{1,2}

¹Department of Biochemistry, Faculty of Biomedical Sciences, Kampala International University - Western Campus, Bushenyi, Western Region, Uganda

²Department of Biochemistry, Faculty of Science, Ebonyi State University, Abakaliki, Ebonyi, Nigeria

³Department of Physiology, School of Medicine, Kabale University, Kabale, Western Region, Uganda

⁴Department of Medical Imaging Sciences, School of Health Sciences, College of Medicine & Health Sciences, University of Rwanda, Butare, Southern Province, Rwanda

⁵Department of Medical Laboratory Science, School of Allied Health Sciences, Kampala International University, Kampala, Central Region, 20000, Uganda

⁶Department of Biochemistry, Faculty of Science, Evangel University Akaeze, Okpoto, Ebonyi, Nigeria

⁷College of Pharmaceutical Sciences, Southwest University College of Pharmaceutical Sciences, Chongqing, Chongqing, China

⁸Department of Anatomy, Faculty of Biomedical Sciences, Kampala International University - Western Campus, Bushenyi, Western Region, Uganda

⁹Department of Publications and Extension, Kampala International University, Kampala, Central Region, Uganda

¹⁰Department of Science Education, Faculty of Education, Kampala International University, Kampala, Central Region, Uganda

V1 First published: 16 Sep 2025, 14:931
<https://doi.org/10.12688/f1000research.169163.1>

Latest published: 16 Sep 2025, 14:931
<https://doi.org/10.12688/f1000research.169163.1>

Open Peer Review

Approval Status AWAITING PEER REVIEW

Any reports and responses or comments on the article can be found at the end of the article.

Abstract

Background

Iron Deficiency Anaemia (IDA) is a major public health challenge in East Africa, particularly affecting children and women. Conventional

therapies are often limited by accessibility and cost, prompting interest in plant-based interventions as affordable and locally available alternatives. This review explores the potential of medicinal plants in managing IDA, focusing on their phytochemical composition, mechanisms of action, and haematological effects.

Methods

A systematic search of PubMed, Scopus, and Web of Science databases was conducted on July 17, 2024. The search strategy utilised Boolean operators and keywords related to plants, IDA management, and East Africa. Studies meeting inclusion criteria were peer-reviewed articles published in English that evaluated plant-based treatments for IDA in preclinical or clinical trials. A total of 30 eligible studies were included after screening and quality assessment following PRISMA guidelines.

Results

The review identified plants such as *Moringa oleifera*, *Phyllanthus emblica*, and *Telfairia occidentalis* as effective in improving haemoglobin levels, red blood cell counts, and other haematological indices. Mechanisms include enhanced iron absorption, antioxidant activity, and stimulation of haematopoiesis. Extraction techniques commonly used were high-performance liquid chromatography (HPLC) and aqueous methods. Key phytochemicals like flavonoids, saponins, and phenolic compounds played critical roles in IDA management. Despite promising outcomes, variations in treatment duration and protocols were noted, highlighting the need for standardisation.

Conclusion

Plant-based therapies offer accessible, cost-effective solutions for managing IDA in East Africa, leveraging local resources and phytochemical properties. However, further pharmacological and toxicological studies are essential to confirm safety and efficacy. This review emphasises the potential of medicinal plants in addressing IDA and calls for the exploration of underutilised plant species and the development of standardised therapeutic protocols.

Keywords

Iron Deficiency Anemia (IDA), Plant-based therapies, Phytochemicals, Hematological indices, East Africa



This article is included in the **Plant Science** gateway.

Corresponding author: Angela Mumbua Musyoka (angela2mumbua@gmail.com)

Author roles: **Siida R:** Conceptualization, Data Curation, Formal Analysis, Methodology, Writing – Original Draft Preparation; **Fasogbon IV:** Conceptualization, Formal Analysis, Methodology, Writing – Original Draft Preparation; **Musyoka AM:** Conceptualization, Data Curation, Resources, Writing – Original Draft Preparation, Writing – Review & Editing; **Nwali BU:** Investigation, Resources, Software, Validation, Visualization; **Etumah SO:** Investigation, Resources, Software, Validation, Visualization; **Okon IA:** Conceptualization, Investigation, Resources, Software, Validation; **Eze ED:** Investigation, Resources, Software, Supervision; **Adam A:** Investigation, Resources, Software, Validation, Visualization; **Mbyemeire H:** Conceptualization, Methodology, Resources, Writing – Original Draft Preparation; **Vandu CD:** Conceptualization, Investigation, Resources, Software, Writing – Original Draft Preparation; **Owembabazi E:** Data Curation, Investigation, Resources, Validation, Visualization; **Okon MB:** Conceptualization, Data Curation, Resources, Visualization, Writing – Original Draft Preparation; **Agu PC:** Resources, Software, Supervision, Writing – Original Draft Preparation; **Ibe UM:** Methodology, Resources, Supervision, Writing – Original Draft Preparation; **Makena W:** Conceptualization, Methodology, Resources, Software, Writing – Original Draft Preparation; **Oviosun A:** Conceptualization, Investigation, Methodology, Writing – Original Draft Preparation; **Etukudo ME:** Conceptualization, Methodology, Visualization, Writing – Original Draft Preparation; **Mujinya R:** Conceptualization, Investigation, Resources, Software; **Dangana RS:** Formal Analysis, Methodology, Resources, Software; **Uti DE:** Investigation, Resources, Software, Validation, Visualization; **Ojiakor VO:** Conceptualization, Methodology, Software, Writing – Original Draft Preparation; **Alum EU:** Data Curation, Investigation, Resources, Supervision, Visualization; **Terkimbi SD:** Methodology, Resources, Software, Validation, Writing – Original Draft Preparation; **Aja L:** Resources, Software, Validation, Visualization; **Jega SA:** Formal Analysis, Resources, Supervision, Writing – Review & Editing; **Paul Chima UO:** Data Curation, Resources, Software, Validation, Visualization; **Mitaki NB:** Investigation, Resources, Software, Validation, Visualization; **Ifie JE:** Resources, Software, Supervision, Writing – Original Draft Preparation; **Mecthilde M:** Investigation, Resources, Software, Validation, Visualization; **Olorunnisola SO:** Formal Analysis, Resources, Supervision, Writing – Original Draft Preparation; **Aja PM:** Conceptualization, Formal Analysis, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2025 Siida R *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Siida R, Fasogbon IV, Musyoka AM *et al.* **Plant-based management of Iron Deficiency Anemia (IDA) in East Africa community: a systematic review [version 1; peer review: awaiting peer review]** F1000Research 2025, 14:931 <https://doi.org/10.12688/f1000research.169163.1>

First published: 16 Sep 2025, 14:931 <https://doi.org/10.12688/f1000research.169163.1>

Introduction

Anemia is a blood disorder characterized by decreased haemoglobin (Hb) concentration. It is defined as haemoglobin levels <13.0 g/dL in men and <12.0 g/dL in women.¹ Other factors that can alter an individual Hb level include ethnicity and physiological status.² Iron deficiency anemia (IDA) is one of the most common causes of anemia. It occurs when there is insufficient iron in the body to produce red blood cells, an important cellular component of blood for the distribution of oxygen to tissues. Sources of iron include protein-rich diets such as fish and meat.³ Studies have shown that IDA accounts for about 50% of anemia in children.⁴ Among other haematological disorders, anemia is one of the biggest causes of disability worldwide and a major public health concern.⁵ Recent statistics have shown a progressive increase in the prevalence of anemia in developed and developing countries in the past decade, especially in sub-Saharan Africa with children accounting for 61.4%.^{6,7}

An estimate from the United Nations revealed that East Africa contributes about 7% of the world population and 21.4% of children in this population suffer from IDA, probably due to the socio-economic status of people living in this region.⁸ According to the 2016 Uganda Demographic and Health Survey (UDHS), the prevalence of iron deficiency anaemia increased from 23% in 2011 to 53% in women of reproductive age and 49% in children aged 6-59 months.⁹ In Kenya, 69% of children under 5 years old bear the brunt of iron deficiency anemia.¹⁰ Sunguya et al.¹¹ reported that the prevalence of IDA among pregnant women in Tanzania has risen to 57% thus the need for immediate intervention.

Several pharmacological interventions have been employed in the management of IDA, especially in East Africa. There is increasing scientific evidence that interventions with medicinal plants have gained global attention in recent years.¹² Plant sources are known to contain phytochemicals with proven pharmacological efficacies in disease conditions. These sources are relatively available and affordable in low and middle-income countries.

Given the discrepancies in recent published studies regarding plant-based management of IDA, we carried out a systematic review. This review aimed to assess different published studies on plant-based management of iron deficiency anaemia (IDA) in the East African community.

Materials and methods

A comprehensive literature search was systematically conducted across the Web of Science (WoS), Scopus, and PubMed databases on 17th July 2024. The search employed the following terms: Plant, management, “iron deficiency anemia”, and “East Africa.” Boolean operators (AND/OR/NOT), alternative terms, and various delimiters such as quotation marks, parentheses, wildcards, and asterisks (*) were utilized to create the search strategy outlined in [Table 1](#), as reported by Fasogbon et al.¹³⁻¹⁵ The search was restricted to peer-reviewed articles published in English. The paper selection process adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 guidelines^{16,17} and followed the inclusion and exclusion criteria presented in [Table 2](#). The risk of bias was assessed as shown in [Figure 5 \(a-b\)](#).

Results

Search results


The Web of Science database search produced 50 articles, Scopus yielded 212, and PubMed identified 30, bringing the total to 292 articles across the three databases. The search results from each database were then exported and imported into Rayyan, a platform specifically designed for systematic review processes, where they were screened based on the inclusion and exclusion criteria.¹⁸ On the platform, 78 duplicate articles were removed. The remaining 214 records were initially screened by evaluating their titles and abstracts, which led to the exclusion of 153 articles. Following a more

Table 1. Search strategies.

Database	Search strategy
Scopus	(Plant*) AND (treat* OR manag* OR therap* OR control* OR ameliorat*) AND (“iron deficiency anemia” OR “iron-deficiency anemia”)
WoS	(Plant*) AND (treat* OR manag* OR therap* OR control* OR ameliorat*) AND (“iron deficiency anemia” OR “iron-deficiency anemia”)
PubMed	((plant*[title/abstract]) AND ((treat*[title/abstract]) OR (manag*[title/abstract]) OR (therap*[title/abstract]) OR (control*[title/abstract]) OR (ameliorat*[title/abstract]))) AND (“iron deficiency anemia”[title/abstract]) OR (“iron-deficiency anemia”[title/abstract])) NOT (review [Publication Type]))

Table 2. Inclusion and exclusion criteria.

Inclusion criteria	Exclusion criteria
• Studies involving the IDA treatment	• Studies not focusing anemia due to iron deficiency as primary or secondary cause
• Treatment must be by plant extract or plant-based product	• Studies focusing on any other type of anemia other than IDA
• Studies must report the impact of the treatment	• Studies that did not describe the impact of plant or plant-based product on IDA
• Studies on either preclinical or clinical trials	• Review articles, editorials, commentaries, book chapters and conference abstracts
• Peer-reviewed articles published in English	• Non-English publications

**Figure 1. PRISMA flow chart for study selection.**

detailed full-text review, an additional 30 articles were excluded for not meeting the inclusion criteria. In the end, 30 articles were included in the study after passing the eligibility criteria and quality assessment (Figure 1).

Table 3 above outlines the mechanisms of action, effects on anemia-related parameters, plants used, plant parts studied, duration of treatment, and geographic origin of the studies.

Mechanisms of action and impact on anemia parameters

Various plants and plant-based combinations have been studied for their roles in increasing hemoglobin levels, red blood cell counts, hematocrit, and other indices of blood health. From the table above, some plants show mechanisms related to enhanced iron absorption or bioavailability, antioxidant action, or stimulation of hematopoiesis. Examples are *Moringa*

Table 3. Mechanisms of action of the plant and parts used in the study.

Mechanism of IDA treatment/ Amelioration	Effect on parameters	Plant	Part of plant	Duration	Country	Reference
Not specified	Increase Hb and pcv, at 2g/dl and 4g/dl, MCV, MCH, MCHC and RCC insignificant at 2g/dl and significant at 4g/dl	Emblic myrobalan	Fruits	60 days	Pakistan	¹⁹
Mixture of honey, dates, and amla (Indian gooseberry)	The combination of honey, dates, and amla boosts hemoglobin levels, increases RBC count, and reduces fatigue, contributing to improved overall vitality	Phyllanthus emblica, Indian gooseberry	Fruits	3 months	India	²⁰
Increased Hemoglobin Levels, increase in Vitamin C levels	significant increase in hemoglobin levels, mean levels of Vitamin I and iron level	Phyllanthus emblica, Cucurbita pepo, Saccharum officinarum	Amla fruit, Jaggeri no part specified, Pumpkin leaves	60 days	India	²¹
Reducing phytate boosts iron bioavailability, and adding Moringa oleifera leaves enhances absorption and nutrition	Not specified	Moringa oleifera	Leaves	12 weeks	Switzerland	²²
Increased iron intake from Hibiscus sabdariffa meals is supported by vitamin C enhancing absorption	Not specified	Microalgae/ Schizochytrium spp	Not specific	xxxx	Ghana	²³
Not specified	Lacto-ovo vegetarians and vegans had lower hemoglobin, erythrocyte, and hematocrit levels, with increased RDW, compared to omnivores	Hibiscus sabdariffa		Not specific	Spain	²⁴
Treatment of IDA in restoration of hematopoiesis by adding decoction of the plant's aerial parts	Effect on erythrocytes (red blood cells), platelets and leukocytes (white blood cells)	Urtica chamaedroides	Leaves, stems, and flowers	34 days	Mexico	²⁵
Through the proliferation of hematopoietic stem cells	Increase RBC, Hemoglobin, Hematocrit decreased μ -RBC, & MRBC), with no significant changes in Platelets	Angelica gigas	Roots and rhizomes	7 days	South Korea	²⁶
Not specified	Haemoglobin, packed cell volume and red blood cells	Sorghum bicolor	Aqueous Moench stem bark	6 weeks	Nigeria	²⁷

Table 3. Continued

Mechanism of IDA treatment/Amelioration	Effect on parameters	Plant	Part of plant	Duration	Country	Reference
Not specified	RBC count, hemoglobin, and hematocrit	Ajwa Date	Fruits	Not specify	Indonesia	²⁸
Iron Supplementation, Improved Iron Storage, and Prevention of Iron Deficiency	RBC count, hemoglobin, and hematocrit	Caulis	Caulis extract	42 days	China	²⁹
Iron Supplement Improved Iron Storage & Prevention of Iron Deficiency	Hemoglobin (Hgb), and Anemia Prevalence	Lentils	Seeds	4 months	Bangladesh	³⁰
Not specified	Hemoglobin, Ferritin, Erythrocyte	Flacourtia indica	Leaf	3 months	India	³¹
Not specified	Hematocrit, MCV, MCH, MCHC, and RDW	Moringa oleifera	Ethanol Leaf extract	3 weeks	Indonesia	³²
Anti-oxidative action of chlorophyll is as a hydrogen donor to break the chain reaction, due to the porphyrin in its chemical structure	CSA increased the levels of Hb, serum iron and serum ferritin	Pearl millet (Pennisetum americanam)	Seeds	30 days	India	³³
Increase in iron bioavailability	Increased Hb,serum iron and serum ferritin levels	Telfairia occidentalis	Leaves and seeds	10 days	United Kingdom	³⁴
Not included	Iron and Hb determined	Mangifera indica L (mango)	Stem bark & leaves	14 days	Nigeria	³⁵
Effects of both iron and phytochemicals of the extract induced boost RBC indices in the iron deficient rats	Increased Hb, concentration, PCV volume; RBC, MCV, & WBC	Plant based diet	Not specified	Not specify	Indonesia	³⁶
Not specified	Increased Hb, concentration, PCV volume; RBC, MCV, & WBC	Angelica sinensis	Dry roots of A. sinensis (Polysaccharide)	14 days	China	³⁷
ASP inhibit hepcidin expression in vivo by stimulating erythropoietin secretion and interrupting two other major pathways of hepcidin regulation	Improves on haematological factors	Punica granatum L (Pomegranate)	Fruits	5 hours	India	³⁸
Increased iron bioavailability in the presence of PJ and Other organic acids, sugars and polyphenols present in PJ might as well have a role in modulation of iron bioavailability	Pomegranate Juice Improves Iron Uptake and Assimilation in Caco-2 Cells. PJ Enhances Iron Uptake and Assimilation in HepG2 Cells	Moon dal washed (MDW), red lentils (RL), chickpeas (CP), green lentils whole (GLW), and yellow split peas (YSP)	Legume seed	Not specified	Canada	³⁹

Table 3. *Continued*

Mechanism of IDA treatment/Amelioration	Effect on parameters	Plant	Part of plant	Duration	Country	Reference
Increased hepcidin mRNA correlates with bioavailable iron, while reduced DM-T1 and TfR1 reflect decreased iron uptake	Iron extracted from defatted microalgae seemed to be effective in alleviating moderate anemia	Defatted Nannochloropsis oceanica (DGM) (defatted microalgae)	Whole plant	10 weeks	United states	40
The macro and micronutrient content in Moringa powder is thought to work synergistically to increase Hb levels in research subjects significantly	Supplement (P2) increased hemoglobin level	Moringa (Moringa oleifera)	Leaves (capsules)	60 days	Indonesia	41
ASP significantly down-regulates JAK1 expression, similar to its effect on phospho-JAK1, while rHPO has a stronger inhibitory effect on JAK1 and its phosphorylation	Increased, RBC, platelets, hematocrit, MCV, MCH & MCHC	Angelica sinensis	Polysaccharides from Angelica sinensis	16 days	China	42
The increase in hemoglobin and erythrocytes in this study is attributed to the presence of iron found in moringa leaves	Increase in hemoglobin and erythrocyte levels	Moringa tree (Moringa oleifera)	Leaves	15 days	Indonesia	43
The mechanism of antioxidative action of chlorophyll is as a hydrogen donor to break the chain reaction, due to the porphyrin in its chemical structure	CSA increased the levels of Hb, serum iron and serum ferritin	Sauropus androgynus (L) Merr (katuk)	Leaves	37 days	Indonesia	44
Increase in iron bioavailability	Increased Hb, serum iron and serum ferritin levels	Low-Phytic Acid Beans and Biofortified Beans	Seeds	14 days (double meal)	Rwanda	45

oleifera, *Angelica gigas*, and *Telfairia occidentalis*. Some other plants, such as *Emblie myrobalan* and *Phyllanthus emblica*, have significantly improved hematological parameters after specific durations.

Plant parts and study locations

Commonly used plant parts include fruits, leaves, roots, seeds, and bark. The studies span multiple countries, including India, Pakistan, Indonesia, China, Nigeria, Rwanda, and Ghana, highlighting the global interest in plant-based anemia management. Some studies explored combinations of plant products, such as honey, dates, and *amla*, which showed synergistic effects in boosting Hb levels and alleviating fatigue.

Duration of intervention and phytochemical contributions

Treatment durations vary widely, from as short as 5 hours (*Punica granatum*) to a few weeks (*Moringa oleifera*), and several months (*lentils*). These variations emphasize the need to standardize treatment protocols for better comparison and effectiveness. From the table, specific phytochemicals, such as chlorophyll, polyphenols, and vitamin C, contribute to increased iron bioavailability and hematological improvements.

Key: Macro-erythrocytes (MRBC), Micro-erythrocytes (μRBC), PCV, packed cell, RBC, red blood, MCV: mean corpuscular volume, Wbc, MCH, mean corpuscular hemoglobin MCHC.

Biochemical techniques used for the extraction of the plants extracts

Figure 2 presents the biochemical techniques used for the extraction of plant extracts studied in the systematic review. The chart highlights that a significant portion of the studies utilized HPLC, which accounts for the largest percentage (17.65%). HPGC, GSMC, and phytate assay follow as the second most common method (11.76%). Common solvents of extraction are water (aqueous), ethanol, methanol, and hexane. The aqueous method is favored due to its simplicity, cost-effectiveness, and suitability for water-soluble polar compounds like flavonoids and polyphenols.⁴⁶ Ethanol is widely used for its ability to extract both polar and non-polar compounds, making it versatile for isolating a range of bioactive phytochemicals.⁴⁷ The choice of extraction method influences the types of bioactive compounds isolated and their subsequent effectiveness in treating iron deficiency anemia. For example, while aqueous extraction might favor compounds with direct iron bioavailability enhancement properties, ethanolic and methanolic methods may yield a wider variety of bioactives, including those with antioxidant and hematopoietic properties. The prevalence of aqueous




Figure 2. Biochemical techniques used for the extraction of the plants extracts.

and ethanolic extractions indicates a preference for techniques that are more feasible in low-resource settings, aligning with the focus on plant-based therapies in developing regions.⁴⁶

Vitamin, mineral, and carbohydrates compositions of the plants used in the management of anemia

Figure 3 illustrates the vitamin and mineral compositions of the plants used in the management of anemia. Iron (36%) is the most prominent mineral highlighted, reflecting its direct role in addressing iron deficiency anemia (IDA). This is expected since IDA arises from insufficient iron required for hemoglobin synthesis. Vitamin C (20%) appears frequently as a vital nutrient. Its role in enhancing iron absorption, particularly from plant-based sources (non-heme iron), highlights its importance in these therapies. Vitamin A (12%) was the second most abundant vitamin. Vitamin A is noted for its antioxidant potential⁴⁸ which contributes to hematopoiesis. Folic acid (4%) is also notable in the chart, as it supports red blood cell formation, highlighting its relevance in anemia management. The chart in Figure 3 highlights the multi-dimensional approach of plant-based management, where the synergy of vitamins and minerals aids not only in addressing iron deficiency but also in improving general nutritional status and hematopoiesis. Notably, plants with high iron content combined with vitamin C are particularly effective, as vitamin C mitigates inhibitors of iron absorption, such as phytates and tannins.⁴⁹

Phytochemical constituents of the plants used in the management of anemia

Figure 4 provides the summary of the phytochemical constituents of the plants used in the management of anemia, highlighting the compounds responsible for therapeutic effects. The phytochemical constituents' range was: Flavonoids>Saponins>Phenolic Compounds> Alkaloids>Others.

Flavonoids (26%) dominate the chart. Flavonoids are known for their antioxidant properties and ability to enhance iron bioavailability. They can chelate free iron, reducing oxidative stress while also protecting erythrocytes from damage.⁵⁰ Saponins (11%) are bioactive compounds that may stimulate erythropoiesis (red blood cell production) and improve gut health, aiding in the absorption of iron and other nutrients.⁴⁹ Phenolic compounds (10%) contribute to antioxidant activity, protecting red blood cells from oxidative damage and supporting the overall health of hematopoietic cells. They also enhance the bioavailability of iron in some situations.⁵¹ Alkaloids (10%) are associated with hematopoietic activity and may stimulate red blood cell production, making them valuable in combating anemia. Others are steroids, terpenes, tannins, and glycosides, which are potentially beneficial in synergistic interactions with the primary phytochemicals. The high presence of flavonoids and phenolic compounds suggests a focus on protecting red blood cells and hematopoietic tissues from oxidative stress, a common challenge in anemia. While tannins can inhibit non-heme iron absorption, their interaction with other phytochemicals (e.g., flavonoids and phenolics) may mitigate this effect. Thus, the combined presence of diverse phytochemicals points to the potential for synergistic effects, where compounds work together to




Figure 3. Vitamin, mineral, and carbohydrates compositions of the plants used in the management of anemia.




Figure 4. Phytochemical constituents of the plants used in the management of anemia. (a) A revised tool to assess the Quality in randomized trials (RoB 2). (b) SYRCLE’s risk of bias tool for animal studies.




Figure 5. (a-b): Risk of bias analysis of the included articles.

enhance therapeutic efficacy, including improving gut health, reducing inflammation, and stimulating red blood cell production.

Discussion

After eligibility screening, only 30 studies from the 292 results of the first literature search were included in the qualitative analysis. Furthermore, there is still a significant amount of information lacking, as seen by the yearly trend of ethnobotanical research on plants used to treat anemia. Since its start, few anemia-related ethnobotanical investigations have been done in East Africa. On the other hand, we have the most ethnobotanical research locally documented in East Africa. Africa has a rich cultural heritage, but there are still more areas to investigate to close the knowledge gap. There is a large probability that other plants utilized for anemia still need to be documented within the African region.

Iron deficiency anemia (IDA) is a condition where the body lacks sufficient iron to produce hemoglobin, a protein in red blood cells that transports oxygen. The pharmacological benefits of bioactive chemicals found in traditionally used plants on iron deficiency anemia have been shown in several research.⁵² Hematological indices, including hemoglobin, hematocrit, red blood cell (RBC), mean corpuscular hemoglobin concentration (MCHC), mean corpuscular volume (MCV), and mean corpuscular hemoglobin (MCH), were used to assess the anti-anemic activity of the plants under study, in particular *Telfairia occidentalis*. In an experimental study conducted on albino rats, it was found that the ethanol leaf extract of *Telfairia occidentalis* induced regeneration of RBC, hematocrit, hemoglobin, MCV, and MCH (Cotoraci et al., 2021). Furthermore, *Telfairia occidentalis* leaf raised PCV levels in a trial on antisickling in sickle cell anemia patients.⁵³

According to another study, the main nutrient for spinal cord hematopoiesis is the particular iron found in *moringa leaves*.⁵⁴ Similarly, *Moringa leaves*' protein and amino acid content function as hematopoietic growth agents. High levels of protein and amino acids, which are essential for controlling blood cell differentiation and proliferation, were reported in its leaves. Additionally, the body absorbs iron more readily due to the vitamin C in *Moringa leaf* extract and other plants.⁵⁵ Its potential effects on the expression of liver hepcidin mRNA expression are suggested by the finding that dietary iron from plants was more effective than ferric citrate in curing iron deficiency in male Wistar rats in a study conducted.^{52,56}

The biochemical techniques used for the extraction of plants in this review revealed that HPLC and GSMC form the bulk of the percentage of techniques used. The choice of a quick and accurate technique to measure the quantity of iron in plant tissues and fluids is a major technological challenge for assessing iron insufficiency in plants. As a precise instrument, HPLC can be used to separate complicated biological samples or synthetic chemicals, analyze organic molecules, confirm medicine authenticity, provide quantitative data, monitor the course of disease therapy, and discover contaminants and degradants in raw materials and final products.⁵⁷ This makes HPLC the most reliable method for the extraction of plant products used in the treatment and management of IDA. Studies have also shown that the iron status of plants under normal growth conditions can be accurately and simply analyzed utilizing an HPLC-based technique.⁵⁸

Among the number of vitamins and minerals identified in works of literature used for this review, Iron constitutes about 36%. IDA can be seen in conditions such as hemorrhage, prolonged menstruation, hookworm infections, and inadequate consumption iron-rich diet.⁵⁹ However, other disease conditions can also prevent iron from being absorbed into the body. Heme (Iron) is a key component of hemoglobin, the oxygen-carrying protein that gives the red blood cell its reddish color. The binding affinity of iron to the protein globin enhances erythropoiesis and the oxygen-carrying capacity in red blood cells. IDA occurs when there is insufficient iron to bind with globin⁶⁰ (Obeagu and Obeagu, 2025). Medicinal plants within East Africa have been proven to increase red cell production due to their high iron content.

Phytochemicals present in these plants reveal their anti-inflammatory and antioxidant properties in the management and treatment of IDA within countries in East Africa. A study has shown that flavonoids can control iron absorption and homeostasis, so they may be useful in the treatment of IDA.⁶¹ Additionally, flavonoids can be utilized to improve red blood cell indices in conjunction with iron supplementation. Also, by controlling the expression and activity of proteins that govern iron uptake and metabolism, flavonoids can have an impact on iron homeostasis and regulate red blood cell production.⁶² Tannin and saponins are active ingredients of secondary metabolites that are recognized to provide several health benefits, including their antioxidant and antibacterial properties.⁶³ These properties can also prevent disease conditions that can result in IDA and improve red blood cell indices.

Conclusion

Thirty screened ethnobotanical articles that documented the use of medicinal plants for the treatment of IDA within the East African region were collected for this systematic review. An increase in traditional medicine research has noted for decades. However, more scientific investigation is required for medicinal plants that are used for treatment and

management of iron deficiency anemia. Ethnobotanical research on IDA is still lacking in some East African regions. The different methods of extraction, minerals and vitamins as well phytochemicals of medicinal plants used to treat IDA in East Africa was demonstrated by this systematic review. To ascertain and confirm the safety and effectiveness of these medicinal plants in treating IDA in the general population, pharmacological and toxicological research is still required. Our current review will direct future researchers to examine unexplored medicinal plants in detail and evaluate their characteristics, focusing on human toxicity and safety.

Data availability statement

Data supporting the findings of this study are available in the Open Science Framework

DOI: <https://doi.org/10.17605/OSF.IO/U95Z8>

This project contains following files:

- PRISMA_2020_checklist.docx
- Revised-Anemia systematic review manuscripts-2025.docx

Data are available under CC0 1.0 Universal license

Acknowledgments

Not applicable.

References

1. Abuaiasha M, Itani H, El Masri R, *et al.*: **Prevalence of iron deficiency (ID) without anemia in the general population presenting to primary care clinics: a cross-sectional study.** *Postgrad. Med.* 2020; **132**(3): 282–287.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Kang W, Barad A, Clark AG, *et al.*: **Ethnic differences in iron status.** *Adv. Nutr.* 2021; **12**(5): 1838–1853.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Sheffield S, Fiorotto ML, Davis TA: **Nutritional importance of animal-sourced foods in a healthy diet.** *Front. Nutr.* 2024; **11**: 1424912.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Obeagu EI, Obeagu GU, Habimana JB: **Iron Deficiency Anaemia in Children.** *International Research in Medical and Health Sciences.* 2023; **6**(4): 26–32.
[Publisher Full Text](#)
5. Fite MB, Assefa N, Mengiste B: **Prevalence and determinants of Anemia among pregnant women in sub-Saharan Africa: a systematic review and Meta-analysis.** *Arch. Public Health.* 2021; **79**: 211–219.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Tirore LL, Areba AS, Habte A, *et al.*: **Prevalence and associated factors of severity levels of anemia among women of reproductive age in sub-Saharan Africa: a multilevel ordinal logistic regression analysis.** *Front. Public Health.* 2024; **11**: 1349174.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Tesema GA, Worku MG, Tessema ZT, *et al.*: **Prevalence and determinants of severity levels of anemia among children aged 6–59 months in sub-Saharan Africa A multilevel ordinal logistic regression analysis.** *PLoS one.* 2021; **16**(4): e0249978.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Hailu BA: **Mapping, trends, and factors associated with anemia among children aged under 5 y in East Africa.** *Nutrition.* 2023; **116**: 112202.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Buyinza F, Muhammad A: **Mothers' education and childhood anaemia in Uganda.** *Dev. South. Afr.* 2024; **41**(2): 427–445.
[Publisher Full Text](#)
10. Kisiangani I, Mbakaya C, Makokha A, *et al.*: **Assessment of iron status among preschool children (6 to 59 months) with and without malaria in Western Province, Kenya.** *Pan Afr. Med. J.* 2015; **21**(1): 62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Sunguya BF, Ge Y, Mlunde L, *et al.*: **High burden of anemia among pregnant women in Tanzania: a call to address its determinants.** *Nutr. J.* 2021; **20**: 65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Zubair S, Sharif A, Abdullah S, *et al.*: **Unlocking the Power of Plants: Herbal Remedies for Anemia.** *International Journal of Trends in Food Science and Technology.* 2023; **1**(1): 1–9.
13. Fasogbon IV, Aja PM, Ondari EN, *et al.*: **UCP-LF and other assay methods for schistosome circulating anodic antigen between 1978 and 2022.** *Biology Methods and Protocols.* 2023; **8**(1).
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Fasogbon IV, Ondari EN, Deusdedit T, *et al.*: **Point-of-care potentials of lateral flow-based field screening for Mycoplasma bovis infections: a literature review.** *Biology Methods and Protocols.* 2024; **9**(1): bpa034.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Fasogbon IV, Yakubu MN, Adam M, *et al.*: **Non-Selelex-Based In-Silico Modeled Aptamers Against Sars-Cov-2 Proteins: A Systematic Review.** *KIU Journal of Health Science.* 2022; **2**(1): 69–79.
16. Page MJ, McKenzie JE, Bossuyt PM, *et al.*: **The PRISMA 2020 statement: An updated guideline for reporting systematic reviews.** *The BMJ.* 2021; **372**.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Mustapha A, Chidiebere O, Victor FI, *et al.*: **Antibacterial activity of Nigerian medicinal plants as panacea for antibiotic resistance: A systematic review.** *Journal of Medicinal Herbs.* 2022; **13**(3): 11–23.
18. Johnson N, Phillips M: **Rayyan for systematic reviews.** *J. Electron. Resour. Librariansh.* 2018; **30**(1): 46–48.
[Publisher Full Text](#)
19. Ritika M, Rizwana KH, Bora B, *et al.*: **Traditional and Underutilized Fruits and Vegetables for Attaining Zero Hunger.** *Food Production, Diversity, and Safety Under Climate Change 2024.* Cham: Springer Nature Switzerland; 2024; pp. 89–111.
20. Chen SY, Chiang IC, Chen YY, *et al.*: **Recent advances in the potential of Phyllanthus emblica L. and its related foods for combating metabolic diseases through methylglyoxal trapping: Phyllanthus emblica L. Attenuates metabolic diseases via MG trapping.** *Food Res. Int.* 2024; **11**: 114907.

21. Habte G, Habte S, Jilo O, *et al.*: **Antimalarial efficacy test of the aqueous crude leaf extract of Coriandrum sativum Linn.: an in vivo multiple model experimental study in mice infected with Plasmodium berghei.** *BMC Complementary Medicine and Therapies.* 2024; **24**(1): 267.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Mawouma S, Hamidou Yaya S, Mbyeya J, *et al.*: **Bioaccessibility and Speciation of Iron from Aqueous Extracts of Moringa oleifera Leaves.** *J. Food Biochem.* 2024; **2024**(1): 9312118.
23. Oyewole AO, Diosady LL: **Evaluating the potential of Hibiscus sabdariffa beverage to address the prevalence of iron deficiency in sub-Saharan Africa.** *LWT.* 2023; **188**: 1154333.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Mavrommatis A, Tsiplakou E, Zerva A, *et al.*: **Microalgae as a sustainable source of antioxidants in animal nutrition, health and livestock development.** *Antioxidants.* 2023; **12**(10): 1882.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Rodolfo VL, Janet RC, Rafaela TA, *et al.*: **Evaluation of hematopoietic and antibacterial activities of Urtica urens L.** *Evaluation.* 2020; **2**: 380–394.
26. Kang M, Park S, Chung Y, *et al.*: **Hematopoietic effects of Angelica gigas Nakai extract on cyclophosphamide-induced myelosuppression.** *Plants.* 2022; **11**(24): 3476.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Fagbohoun L, Nonvidé GC, Orou AS, *et al.*: **Anti-anaemic activity and potential toxicity of extracts of four tinctorial plants used in the treatment of anemia in Benin: Gossypium barbadense, Sorghum bicolor, Hibiscus sabdariffa and Justicia secunda.** *Am. J. Plant Sci.* 2022; **13**(12): 1460–1477.
[Publisher Full Text](#)
28. Xu J, Liu J, Li B, *et al.*: **Comparison of blood tonic efficacy and chemical constituents of Kadsura interior AC Smith and its closely related species.** *Chin. Med.* 2022; **17**(1): 14.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Soltan SS: **The protective effect of soybean, sesame, lentils, pumpkin seeds and molasses on iron deficiency anemia in rats.** *World Appl. Sci. J.* 2013; **23**(6): 795–807.
30. Swati M, Nath SG, Yatendra K, *et al.*: **Phytochemical analysis and free-radical scavenging activity of Flacourtia indica (Burm. F.).** *Merry Journal of Pharmaceutical Research.* 2009; **8**(2): 81–84.
31. Perera C, Galappatti DI, Thrimavithana AM, *et al.*: **Effect of Moringa oleifera on haematological parameters: a systematic review.** *J. Coll. Commun. Sri Lanka.* 2022; **28**(2): 602.
[Publisher Full Text](#)
32. Alasa MC: **Nutritional Assessment of Panicum maximum (Jacq.) ensiled with two cultivars of Lablab purpureus (Lablab purpureus L.) for West African Dwarf Ram.** 2014. (Doctoral dissertation). [Accessed on March 7, 2025].
[Reference Source](#)
33. Okokon JE, Osigwe CC, Andrew UE, *et al.*: **Effect of Telfairia occidentalis Stem Extract and Fractions on Parasitaemia, Oxidative Stress Markers, Lipid Profile, Hematological Parameters, Liver Function Indices, and Liver Histology.** *Asian Journal of Biochemistry, Genetics and Molecular Medicine.* 2025; **17**(1): 42–57.
34. Asanga EE, Okoroiwu HU, Umoh EA, *et al.*: **Antiplasmodial Profiling of Mangifera indica's Herbal Formulation and Its Ability to Ameliorate Hematological Derangements in Plasmodium berghei-Infected Mice.** *Nat. Prod. Commun.* 2023; **18**(12): 1934578X231215937.
[Publisher Full Text](#)
35. Rahfiludin MZ, Arso SP, Joko T, *et al.*: **Plant-based Diet and Iron Deficiency Anemia in Sundanese Adolescent Girls at Islamic Boarding Schools in Indonesia.** *Journal of Nutrition and Metabolism.* 2021; **2021**(1): 1–7.
[Publisher Full Text](#)
36. Younas F, Aslam B, Muhammad F, *et al.*: **Haematopoietic effects of Angelica sinensis root cap polysaccharides against lisinopril-induced anaemia in albino rats.** *Pharm. Biol.* 2017; **55**(1): 108–113.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Niewiadomska J, Kumiega E, Póciennik M, *et al.*: **Effects of Punica granatum L. peel extract supplementation on body weight, cardiac function, and haematological and biochemical parameters in an animal model of metabolic syndrome.** *Journal of Vet. Res.* 2023; **67**(2): 219–232.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Saeed M, Naveed M, Bibi J, *et al.*: **The promising pharmacological effects and therapeutic/medicinal applications of Punica granatum L. (Pomegranate) as a functional food in humans and animals.** *Recent Patents Inflamm. Allergy Drug Discov.* 2018; **12**(1): 24–38.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Gesinde FA, Udechukwu MC, Aluko RE: **Structural and functional characterization of legume seed ferritin concentrates.** *J. Food Biochem.* 2018; **42**(3): e12498.
40. Manor ML, Kim J, Derksen TJ, *et al.*: **Defatted microalgae serve as a dual dietary source of highly bioavailable iron and protein in an anemic pig model.** *Algal Res.* 2017; **26**: 409–414.
41. Otitoju O, Nwamarah JU, Otitoju GT, *et al.*: **Effect of Moringa oleifera aqueous leaf extract on some haematological indices in Wistar rats.** *Chemical and Process Engineering Research.* 2014; **18**: 26–30.
42. Chang CW, Chen YM, Hsu YJ, *et al.*: **Protective effects of the roots of Angelica sinensis on strenuous exercise-induced sports anemia in rats.** *J. Ethnopharmacol.* 2017; **193**: 169–178.
[Publisher Full Text](#)
43. Suzana D, Suyatna FD, Andrajati R, *et al.*: **Effect of Moringa oleifera leaves extract against hematology and blood biochemical value of patients with iron deficiency anemia.** *J. Young Pharm.* 2017; **9**(1s): S79.
44. Suparmi S, Fasitasari M, Martosupono M, *et al.*: **Hypoglycemic and antianemia effects of chlorophyll from sauropus androgynus (L) merr leaves in rats.** *Pharm. J.* 2021; **13**(4): 924–932.
[Publisher Full Text](#)
45. Philipo M: **Iron and zinc genetic biofortification of yellow common bean (phaseolus vulgaris L.) genotypes in Tanzania.** NM-AIST; 2021. [Accessed on March 7, 2025]. (Doctoral dissertation).
[Reference Source](#)
46. Bitwell C, Indra SS, Luke C, *et al.*: **A review of modern and conventional extraction techniques and their applications for extracting phytochemicals from plants.** *Scientific African.* 2023; **19**: e01585.
[Publisher Full Text](#)
47. Awad AM, Kumar P, Ismail-Fitry MR, *et al.*: **Green Extraction of Bioactive Compounds from Plant Biomass and Their Application in Meat as Natural Antioxidant.** *Antioxidants (Basel, Switzerland).* 2021; **10**(9): 1465.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Alum EU, Aja W, Ugwu OPC, *et al.*: **Assessment of Vitamin Composition of Ethanol Leaf and Seed Extracts of Datura Stramonium.** *Avicenna Journal of Medical Biochemistry.* 2023; **11**(1): 92–97.
[Publisher Full Text](#)
49. Piskin E, Cianciosi D, Gulec S, *et al.*: **Iron Absorption: Factors, Limitations, and Improvement Methods.** *ACS Omega.* 2022; **7**(24): 20441–20456.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Wang X, Li Y, Han L, *et al.*: **Role of Flavonoids in the Treatment of Iron Overload.** *Front. Cell Dev. Biol.* 2021; **9**: 685364.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Ousaaid D, Ghouzi AE, Laaroussi H, *et al.*: **Anti-Anemic Effect of Antioxidant-Rich Apple Vinegar against Phenylhydrazine-Induced Hemolytic Anemia in Rats.** *Life (Basel, Switzerland).* 2022; **12**(2): 239.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Cotoraci C, Ciceu A, Sasu A, *et al.*: **Natural antioxidants in anemia treatment.** *Int. J. Mol. Sci.* 2021; **22**(4): 1883.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Cyril-Olutayo MC, Agbedahunsi JM, Akinola NO: **Studies on the effect of a nutritious vegetable, Telfairia occidentalis, on HbSs blood.** *J. Tradit. Complement. Med.* 2019; **9**(2): 156–162.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
54. Bell V, Varzakas T, Psaltopoulou T, *et al.*: **Sickle cell disease update: new treatments and challenging nutritional interventions.** *Nutrients.* 2024; **16**(2): 258.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. Rahmawati R, Zakiah N, Frengki F: **Study of the Potential of the Active Compounds of Moringa oleifera Leaves in Increasing the Intake of Fe+ 3 in Pregnant women through Docking and Molecular Dynamics Methods on the Enzyme" Human Duodenal Cytochrome B"(Dcytb).** *Research Journal of Pharmacy and Technology.* 2024; **17**(10): 4785–4795.
[Publisher Full Text](#)
56. Magtalas MC, Balbin PT, Cruz EC, *et al.*: **Ethnomedicinal plants used for the prevention and treatment of anemia in the Philippines: a systematic review.** *Tropical Medicine and Health.* 2023; **51**(1): 27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. Chen S, Li Z, Zhang S, *et al.*: **Emerging biotechnology applications in natural product and synthetic pharmaceutical analyses.** *Acta Pharm. Sin. B.* 2022; **12**(11): 4075–4097.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
58. Menezes FA, Oliveira JG, Guimarães AO: **Electron Paramagnetic Resonance Applied to Free Radicals and Reactive Oxygen Species Detection in Plant Systems.** *Appl. Magn. Reson.* 2024; **55**(4): 335–355.
[Publisher Full Text](#)
59. Iolascon A, Andolfo I, Russo R, *et al.*: **Recommendations for diagnosis, treatment, and prevention of iron deficiency and**

- iron deficiency anemia.** *Hemasphere.* 2024; **8**(7): e108.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
60. Obeagu EI, Obeagu GU: **Anemia and Cerebrovascular disease: pathophysiological insights and clinical implications.** *Ann. Med. Surg.* 2025; **87**: 3254–3267.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
61. Armi Y, Felina M, Hikmah Z, *et al.*: **The Hematopoietic Potential of Tamarillo (*Cyphomandra betacea*) and Pitaya (*Hylocereus spp.*) Juices in Anemia Management: A Randomized Controlled Trial in Bukittinggi, West Sumatra, Indonesia.** *Eureka Herba Indonesia.* 2024; **5**(3): 492–502.
[PubMed Abstract](#) | [Publisher Full Text](#)
62. Liu J, Deng L, Qu L, *et al.*: **Herbal medicines provide regulation against iron overload in cardiovascular diseases: Informing future applications.** *J. Ethnopharmacol.* 2024; **326**: 117941.
[PubMed Abstract](#) | [Publisher Full Text](#)
63. Hagaggi NSA, Abdul-Raouf UM, Radwan TA: **Variation of antibacterial and antioxidant secondary metabolites and volatiles in leaf and callus extracts of Phulai (*Acacia Modesta* Wall.).** *BMC Plant Biol.* 2024; **24**(1): 93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research