

<https://doi.org/10.59298/NIJSES/2026/71.2024>

# Case Studies of Malaria Surges after Extreme Climate Events in East and West Africa

Nagawa Jackline Irene

Department of Clinical Medicine and Dentistry Kampala International University Uganda  
Email: irene.nagawa@studwc.kiu.ac.ug

## ABSTRACT

Malaria remains one of the most prevalent and deadly infectious diseases in sub-Saharan Africa, where it continues to burden public health systems despite global efforts to control and eliminate it. Climate change has exacerbated malaria transmission, with extreme climate events such as floods, droughts, and temperature fluctuations significantly influencing the spread of the disease. This review examines case studies from East and West Africa, regions where the impact of climate change on malaria transmission has been particularly pronounced. The study highlights the correlations between extreme climate events and surges in malaria incidence, focusing on the role of climate variables like temperature, rainfall, and humidity in facilitating mosquito breeding and parasite development. East Africa, with its varied topography, and West Africa, with its tropical climate, have both experienced notable malaria outbreaks linked to climate anomalies. These case studies underscore the complex relationship between climate variability and malaria transmission, revealing the need for localized, evidence-based strategies to address these challenges. The review emphasizes the importance of integrating climate-sensitive malaria control strategies into public health systems, which include improving weather forecasting, utilizing climate data for early warning systems, and enhancing surveillance efforts. Given the increasing frequency of extreme weather events due to climate change, this research advocates for adaptive public health responses that can mitigate the impact of malaria surges and safeguard vulnerable populations in Africa.

**Keywords:** Malaria, Climate Change, Extreme Weather Events, Africa, Surveillance, Public Health.

## INTRODUCTION

Malaria remains one of the most prevalent and deadly infectious diseases in sub-Saharan Africa, despite extensive global efforts to control and eliminate it. The disease, caused by *Plasmodium* parasites transmitted through the bites of infected Anopheles mosquitoes, continues to place a heavy burden on the region's health systems [1]. According to the World Health Organization (WHO), sub-Saharan Africa accounts for approximately 90% of the global malaria burden, with millions of cases reported annually, resulting in hundreds of thousands of deaths, predominantly among children under five years of age. Malaria transmission is strongly influenced by environmental factors, especially climate conditions [2]. In recent decades, the role of climate change in altering malaria transmission dynamics has gained increasing attention, as the region experiences a rise in extreme climate events, including floods, droughts, and temperature fluctuations. The relationship between climate and malaria transmission is complex, involving a multitude of factors such as temperature, rainfall, humidity, and seasonal variations [3]. While certain climatic conditions are conducive to the proliferation of Anopheles mosquitoes and the development of the *Plasmodium* parasite, extreme climate events, especially those associated with global warming, can exacerbate malaria transmission by creating favorable breeding grounds for mosquitoes, disturbing traditional ecological balances, and

altering the habitats of both the mosquito vectors and the parasite [4]. Understanding the links between these extreme climate events and malaria surges is essential for improving the prediction, prevention, and control of malaria in sub-Saharan Africa. This review examines case studies from East and West Africa, regions where the impact of climate change on malaria has been particularly pronounced. These case studies provide valuable insights into the correlations between extreme climate events and the frequency of malaria outbreaks [5]. By highlighting these patterns, the review aims to contribute to the growing body of research on climate change and infectious diseases, with a focus on malaria as a case study for understanding the broader implications of climate-induced health risks in vulnerable regions [6].

The influence of climate on malaria transmission is not a new concept, with studies dating back several decades. However, recent shifts in global climate patterns, driven by human activities such as deforestation, industrial emissions, and land-use changes, have led to more frequent and severe extreme weather events, including floods, heatwaves, and prolonged droughts. These events not only disrupt the environment but also amplify existing vulnerabilities in health systems, particularly in resource-poor regions like sub-Saharan Africa [7]. Malaria transmission is sensitive to climatic variables. For instance, temperature directly affects the development rate of *Plasmodium* parasites within mosquitoes, with higher temperatures accelerating parasite development, thus shortening the incubation period. This means that mosquitoes in warmer climates have the potential to transmit the disease more quickly. Rainfall, on the other hand, creates breeding sites for mosquitoes, as stagnant water in pools, puddles, and swamps provides ideal conditions for mosquito larvae. Changes in rainfall patterns, therefore, have significant implications for malaria transmission. Similarly, periods of drought may reduce the availability of breeding sites, but in some cases, drought followed by heavy rains can create a sudden spike in mosquito populations, leading to malaria surges [8].

In the context of climate change, these interactions between climate and malaria dynamics are expected to become more unpredictable. Some regions may experience longer dry seasons that disrupt mosquito breeding cycles, while others may see an increase in the frequency of intense rainfall events, further complicating malaria control efforts. This underscores the need for localized, evidence-based strategies that consider the unique climatic and ecological conditions of each region [9]. This study focuses specifically on the East and West African regions, both of which have experienced significant climate shifts over recent decades. East Africa, which includes countries such as Kenya, Tanzania, and Uganda, is characterized by varied topography, with highland areas offering potential refuges for malaria transmission in a changing climate. West Africa, home to countries like Ghana, Nigeria, and Mali, has a more uniform tropical climate, but is also prone to significant fluctuations in rainfall and temperature [10]. Both regions provide valuable case studies for examining the direct and indirect effects of extreme climate events on malaria transmission and the broader public health implications.

Malaria remains a major public health challenge in sub-Saharan Africa, and its incidence is closely tied to environmental conditions. With climate change intensifying the frequency and severity of extreme weather events, there is growing concern about their potential impact on malaria transmission. There is a lack of comprehensive studies that specifically address the relationship between extreme climate events, such as floods, droughts, and temperature shifts, and malaria surges in the context of East and West Africa [11]. While the links between climate and malaria are widely acknowledged, the specific mechanisms by which extreme events influence malaria outbreaks are not well understood. Moreover, many health systems in sub-Saharan Africa are already under strain due to limited resources, inadequate infrastructure, and ongoing disease burdens. As the impacts of climate change become more pronounced, health systems in these regions will face increasing pressure to respond to changing malaria patterns [12]. Failure to understand the nuances of climate-malaria interactions could undermine efforts to control malaria and mitigate its effects, exacerbating the disease burden in an already vulnerable region. This study aims to explore the impact of extreme climate events on malaria transmission and outbreaks in East and West Africa. By analyzing case studies from these regions, it seeks to identify specific climate events, such as temperature fluctuations, heavy rainfall, floods, and droughts, that have triggered significant surges in malaria. Additionally, the study aims to examine the ecological and epidemiological mechanisms through which these extreme climate events influence malaria transmission dynamics. Understanding how factors like changes in temperature and rainfall patterns affect malaria transmission is essential for pinpointing the pathways through which climate change exacerbates the disease. The study will also evaluate the effectiveness of public health responses to malaria outbreaks that follow extreme climate events, assessing how well interventions have mitigated the impact of these outbreaks. Based on the findings, the study will provide recommendations for adapting malaria control strategies to better address the challenges posed by climate change. The significance of this research lies in its ability to enhance the understanding of how climate variability influences malaria patterns in sub-Saharan Africa. By providing valuable insights into the relationship between climate events and malaria, the study will inform targeted and adaptive strategies for malaria control, helping policymakers and health professionals prepare for future outbreaks in a changing climate.

### **The Role of Climate Events in Malaria Transmission**

Climate events play a crucial role in the transmission dynamics of malaria by influencing the various stages of the Plasmodium parasite's lifecycle and the population of its mosquito vectors. Temperature, rainfall, and humidity are particularly significant climate variables that directly affect the rate of malaria transmission. Warmer temperatures, for example, accelerate the development of the Plasmodium parasite within the mosquito host, leading to faster transmission cycles and a higher likelihood of infection [13]. Similarly, increased rainfall provides abundant breeding sites for mosquitoes, particularly in stagnant water pools, which facilitates the growth of mosquito populations. As a result, malaria transmission tends to increase in regions experiencing higher rainfall and warmer temperatures. On the other hand, extreme droughts or extended periods of dry weather can temporarily reduce mosquito populations due to the scarcity of water sources for breeding. However, such droughts are often followed by rapid surges in transmission once the climate becomes favorable again, such as after a period of heavy rainfall. This cyclical pattern highlights the sensitivity of malaria transmission to climate variability and underscores the complex relationship between climate events and the seasonal and geographical patterns of malaria outbreaks [14]. Understanding these links is essential for predicting and controlling malaria transmission in the face of changing climate conditions.

#### **Case Studies**

**East Africa:** East Africa, with its diverse climatic zones, has experienced several malaria surges linked to extreme climate events. A notable example is the 1997-1998 El Niño event in Kenya, which caused widespread flooding and created ideal breeding conditions for Anopheles mosquitoes, the primary vectors of malaria. The heavy rainfall led to stagnant water pools, fostering mosquito proliferation and a dramatic increase in malaria cases [15]. Similarly, Uganda faced severe flooding during the same climatic anomaly, which significantly impacted both malaria incidence and mortality. Health authorities reported a surge in hospital admissions, especially in lowland areas where water stagnation was prevalent, highlighting the direct link between flooding and malaria outbreaks. In Ethiopia, the 2015 flooding event also triggered a surge in malaria cases in regions that had previously experienced low incidence [16]. The floodwaters created new mosquito breeding sites, which facilitated the spread of the malaria parasite to populations that had not encountered malaria for decades. This pattern underscores the complex relationship between extreme weather events and malaria transmission. Climate change-driven events like these do not just exacerbate existing malaria hotspots but also introduce new risks, particularly in regions with historically low malaria transmission, further complicating efforts to control the disease in the region [17].

**West Africa:** West Africa has consistently demonstrated a strong correlation between climate extremes, particularly droughts and floods, and surges in malaria cases. The Sahel region, in particular, provides a notable example. During the 2010-2011 drought, the subsequent heavy rains led to a sharp increase in malaria transmission across countries like Niger, Chad, and Mali. The drought had initially dried up many seasonal pools, which are essential mosquito breeding sites. However, once the rains returned, they replenished these pools, creating ideal conditions for mosquito larvae to thrive [18]. This sudden influx of breeding grounds contributed to a significant uptick in malaria cases, as the mosquitoes that breed in stagnant water transmitted the disease to local populations. A particularly striking case occurred in Ghana's coastal regions after the devastating 2007 floods. The combination of rising humidity and widespread flooding created optimal conditions for the malaria vector, Anopheles mosquitoes, to flourish. Local health systems struggled to cope with the resulting surge in malaria cases. In some regions, health facilities reported up to a 40% increase in malaria-related consultations, overwhelming healthcare providers. These case studies underscore the growing threat that climate extremes pose to malaria transmission in West Africa, highlighting the urgent need for adaptive public health strategies to mitigate the impact of such climate events [19].

#### **Implications for Public Health and Climate Adaptation**

The growing body of research and case studies on malaria transmission highlights the urgent need for an integrated approach to malaria control that effectively incorporates climate predictions and early warning systems. Climate change is increasingly recognized as a key driver of shifts in malaria transmission patterns, influencing both the frequency and severity of the disease. Rising temperatures, fluctuating rainfall patterns, and more frequent extreme weather events, such as floods and droughts, have a direct impact on the habitat of malaria-carrying mosquitoes, thereby altering transmission dynamics [20]. These changing patterns make it difficult for health systems to anticipate and respond to malaria outbreaks in a timely and effective manner.

Health systems must, therefore, be prepared to respond not only to the immediate surge of malaria cases following extreme weather events but also to the long-term shifts in transmission patterns due to broader climatic changes. Traditional preventive measures, such as the distribution of insecticide-treated nets (ITNs) and indoor residual spraying (IRS), remain essential components of malaria control strategies [21]. However, these measures need to be augmented with climate-sensitive approaches that allow for more proactive and anticipatory responses [22]. For example, improving weather forecasting and utilizing climate data to predict malaria risks in specific regions can enable health authorities to adjust their interventions in real time. In addition, surveillance systems should be

strengthened to track environmental and climatic factors that influence mosquito populations and disease transmission.

A key component of effective climate adaptation is fostering collaboration between meteorological agencies, health ministries, and local communities. Meteorological agencies can provide early warnings and climate projections, while health ministries must ensure that this information is used to adjust control measures appropriately. Communities, as frontline stakeholders, need to be engaged and educated about the links between climate change and malaria risks [23]. This collaborative approach not only enhances the effectiveness of malaria control but also builds resilience in vulnerable populations facing the compounded threat of climate change and infectious diseases.

### CONCLUSION

The case studies from East and West Africa highlight the critical link between climate change and malaria transmission, revealing how extreme climate events, such as floods, droughts, and temperature fluctuations, significantly impact malaria incidence. These climate anomalies create environments conducive to the proliferation of malaria vectors, leading to higher transmission rates and increasing the burden on already strained public health systems. The evidence presented emphasizes the urgent need for malaria control programs to integrate climate resilience into their strategies. This includes developing adaptive approaches that can respond to the evolving challenges posed by climate variability. As climate change continues to alter disease transmission patterns, it becomes essential for malaria-endemic countries to craft comprehensive, climate-smart public health strategies. These strategies must combine robust climate science with effective public health interventions, ensuring preparedness and responsiveness to malaria outbreaks. Strengthening such integrated approaches will be crucial for mitigating the effects of malaria surges and safeguarding public health in the face of climate change.

### REFERENCES

1. Mezieobi, K.C., Alum, E.U., Ugwu, O.P.C., Uti, D.E., Alum, B.N., Egba, S.I., Ewah, C.M. Economic burden of malaria on developing countries: A mini review. *Control.* 30 (2025), e00435. <https://doi.org/10.1016/j.parepi.2025.e00435>
2. Malaria | WHO | Regional Office for Africa, <https://www.afro.who.int/health-topics/malaria>
3. Ugwu O P C , Nwodo O F C, Joshua P E, Odo C E, Bawa A, Ossai E. C, Adonu C. C (2019). Anti-malaria and hematological analyses of ethanol leaf extract of *Moringa oleifera* on malaria-infected mice. *International Journal of Pharmacy and Biological Science* 3(1) 360-371.
4. Rubuga, F.K., Ahmed, A., Siddig, E., Sera, F., Moirano, G., Aimable, M., et al.: Potential impact of climatic factors on malaria in Rwanda between 2012 and 2021: a time-series analysis. *Malaria Journal.* 23, 274 (2024). <https://doi.org/10.1186/s12936-024-05097-5>
5. Egwu, C. O., Alope, C., Chukwu, J., Agwu, A., Alum, E., Tsamesidis, I, et al. A world free of malaria: It is time for Africa to actively champion and take leadership of elimination and eradication strategies. *Afr Health Sci.* 2022 Dec;22(4):627-640. doi: 10.4314/ahs.v22i4.68.
6. Rubuga, F.K., Ahmed, A., Siddig, E., Sera, F., Moirano, G., Aimable, M., et al.: Potential impact of climatic factors on malaria in Rwanda between 2012 and 2021: a time-series analysis. *Malaria Journal.* 23, 274 (2024). <https://doi.org/10.1186/s12936-024-05097-5>
7. Kungu, E., Inyangat, R., Ugwu, O.P.C., Alum, E. U. (2023). Exploration of medicinal plants used in the management of malaria in Uganda. *Newport international journal of research in medical sciences* 4(1):101-108.
8. Nabatanzi, M., Ntono, V., Kamulegeya, J., Kwesiga, B., Bulage, L., Lubwama, B., et al.: Malaria outbreak facilitated by increased mosquito breeding sites near houses and cessation of indoor residual spraying, Kole district, Uganda, January-June 2019. *BMC Public Health.* 22, 1898 (2022). <https://doi.org/10.1186/s12889-022-14245-y>
9. Alum, E. U., Tufail, T., Agu, P. C., Akinloye, D. I., Obaroh, I. O. (2024). Malaria pervasiveness in Sub-Saharan Africa: Overcoming the scuffle. *Medicine*, 103(49), e40241. doi: 10.1097/MD.00000000000040241. PMID: 39654176
10. Taye, M.T., Dyer, E.: Hydrologic Extremes in a Changing Climate: a Review of Extremes in East Africa. *Current Climate Change Reports.* 10, 1–11 (2024). <https://doi.org/10.1007/s40641-024-00193-9>
11. Obeagu E. I, Obeagu G, U, Egba S. I, Emeka-Obi O. R, (2023) Combatting Anaemia in Paediatric Malaria: Effective management strategies *Int. J. Curr. Res. Med. Sci.* 9(11): 1-7
12. Rubuga, F.K., Ahmed, A., Siddig, E., Sera, F., Moirano, G., Aimable, M., et al.: Potential impact of climatic factors on malaria in Rwanda between 2012 and 2021: a time-series analysis. *Malaria Journal.* 23, 274 (2024). <https://doi.org/10.1186/s12936-024-05097-5>
13. Emmanuel I. N., Ani. O. C., Ugwu F. J., Egba S. I., Aguzie I. O., Okeke O. P., Dialoke C. E., Asogwa L. O., Odo S. I. (2020) Malaria Prevalence in Rice Farm Settlements South East Nigeria. *IJTDH*, 41(9): 64-74

14. Nosrat, C., Altamirano, J., Anyamba, A., Caldwell, J.M., Damoah, R., Mutuku, F., Ndenga, B., LaBeaud, A.D.: Impact of recent climate extremes on mosquito-borne disease transmission in Kenya. *PLoS Negl Trop Dis.* 15, e0009182 (2021). <https://doi.org/10.1371/journal.pntd.0009182>
15. Lindsay, S.W., Bødker, R., Malima, R., Msangeni, H.A., Kisinza, W.: Effect of 1997–98 El Niño on highland malaria in Tanzania.
16. Onohuean H, Uti DE, Echegu D A. et al Mitigation of Malaria in Sub-Saharan Africa through Vaccination: A Budding Road Map for Global Malaria Eradication. "Ethiopian Journal of Health Sciences. 2025;35(3):205-217. doi: 10.4314/ejhs.v35i3.9. PMID: 40717722; PMCID: PMC12287706.
17. Coalson, J.E., Anderson, E.J., Santos, E.M., Madera Garcia, V., Romine, J.K., Luzingu, J.K., et al.: The Complex Epidemiological Relationship between Flooding Events and Human Outbreaks of Mosquito-Borne Diseases: A Scoping Review. *Environ Health Perspect.* 129, 096002 (2021). <https://doi.org/10.1289/EHP8887>
18. Armin R., Michele R. R., (2013). Malaria during pregnancy with parasite sequestration in the villous chamber. *Blood, the Journal of the American Society of Hematology*, 121, (12), 2173-2173. <https://doi.org/10.1182/blood-2012-10-465096>
19. Duque, C., Lubinda, M., Matoba, J., Sing'anga, C., Stevenson, J., Shields, T., Shiff, C.J.: Impact of aerial humidity on seasonal malaria: an ecological study in Zambia. *Malaria Journal.* 21, 325 (2022). <https://doi.org/10.1186/s12936-022-04345-w>
20. Ugwu, O.P.C., Egba, S.I., Uti, D.E., Alum, B.N. Climate Variability and Malaria Transmission: Unraveling the Complex Relationship. *INOSR Scientific Research.* 2024; 11(2):16-22. <https://doi.org/10.59298/INOSRSR/2024/1.1.21622>
21. Fatima, S.H., Zaidi, F., Rafiq, J., Bhandari, D., Ali, A., Bi, P.: Impact of temperatures on malaria incidence in vulnerable regions of Pakistan: empirical evidence and future projections | *Epidemiology & Infection.* Cambridge Core. <https://doi.org/10.1017/S0950268825000111>
22. Ebele J. I, Emeka E. N., Nnenna C. A., Ignatius C. M., Emeka G. A. (2010). Malaria parasitaemia: effect on serum sodium and potassium levels. *Biol Med*, 2, 2, 20-25.
23. Pinto, J., Cossa, N., Ferrari, M., Coffey, P.S., Picolo, M., Marrufo, T., Cardoso, A.P.: Integration and use of climate data by the national health system in Mozambique. *The Journal of Climate Change and Health.* 22, 100368 (2025). <https://doi.org/10.1016/j.jocl.2024.100368>

**CITE AS: Nagawa Jackline Irene. (2026). Case Studies of Malaria Surges after Extreme Climate Events in East and West Africa. NEWPORT INTERNATIONAL JOURNAL OF SCIENTIFIC AND EXPERIMENTAL SCIENCES, 7(1):20-24. <https://doi.org/10.59298/NLJSES/2026/71.2024>**