

<https://doi.org/10.59298/NIJPP/2026/721418>

The Effectiveness of Borehole Water Systems in Reducing Diarrhea in Rural West Africa: A Review

Mugisha Emmanuel K.

Faculty of Science and Technology Kampala International University Uganda

ABSTRACT

Access to clean drinking water is essential for public health, particularly in rural areas of West Africa, where waterborne diseases such as diarrhea remain a significant threat. Borehole water systems have become a key solution to providing safe drinking water in these communities. This review examines the effectiveness of borehole water systems in reducing the incidence of diarrhea, a major health issue in rural West Africa. The study analyzes existing literature and evidence, focusing on the role of boreholes in improving water quality and public health outcomes, particularly in children under five. The review highlights both successes and challenges, including the importance of community engagement, proper maintenance, water quality monitoring, and hygiene education. While boreholes have proven to reduce diarrheal diseases in some areas, issues such as system sustainability, contamination risks, and inequitable access remain significant obstacles. The findings suggest that for borehole systems to be effective in the long term, improvements in maintenance, monitoring, and community involvement are essential. This review provides valuable insights for policymakers, NGOs, and development practitioners working to improve water access and public health in similar settings.

Keywords Borehole water systems, waterborne diseases, diarrhea, rural West Africa, public health, water quality.

INTRODUCTION

Access to clean and safe drinking water is a fundamental determinant of public health, especially in regions where infectious diseases are prevalent. In rural West Africa, where waterborne diseases continue to pose a significant public health threat, ensuring the availability of safe water is a critical step in addressing the region's health challenges [1]. Among the various public health interventions aimed at reducing waterborne diseases, the installation of borehole water systems has gained prominence in rural communities. These systems are designed to provide a reliable and clean water source, potentially reducing the incidence of waterborne diseases, including diarrhea. Diarrhea, primarily caused by the consumption of contaminated water, is a major cause of morbidity and mortality in rural West Africa [2]. The impact is especially severe for young children, who are more vulnerable to dehydration and malnutrition resulting from diarrhea. The persistent challenge of unsafe drinking water in rural communities calls for a deeper understanding of the effectiveness of borehole water systems in addressing this issue. This review seeks to assess the role of boreholes in reducing the incidence of diarrhea in these areas, identifying both successes and challenges associated with their implementation [3].

West Africa remains one of the regions with the highest burden of waterborne diseases globally. According to the World Health Organization (WHO), millions of people in rural West Africa lack access to safe drinking water, and many rely on unprotected surface water sources, such as rivers, ponds, and streams [4]. These water sources are highly susceptible to contamination from various pathogens, including bacteria, viruses, and parasites, all of which contribute to the spread of diarrhea, cholera, and other waterborne diseases. The problem of unsafe drinking water is compounded by inadequate sanitation facilities and poor hygiene practices. In rural communities, where infrastructure and resources are limited, the lack of access to basic sanitation services and health education often exacerbates the spread of waterborne diseases [5]. Diarrheal diseases, caused by consuming contaminated water

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

and poor sanitation, are a leading cause of death in the region, particularly among vulnerable populations such as children under five and the elderly. The widespread prevalence of diarrhea and other gastrointestinal diseases has significant socio-economic implications. Beyond the direct health impacts, the cost of treating waterborne diseases and the loss of productivity due to illness place a substantial burden on households and healthcare systems in the region [6]. As a result, addressing the root causes of waterborne diseases, poor water quality and inadequate sanitation is vital for improving public health outcomes and reducing preventable deaths in rural West Africa.

The prevalence of waterborne diseases, particularly diarrhea, remains alarmingly high in rural West Africa. Despite significant efforts to improve water access and sanitation, many communities continue to rely on unsafe water sources, contributing to the persistence of diarrheal diseases. Borehole water systems have been implemented as one of the key solutions to improve access to clean drinking water [7]. However, there is limited evidence on the long-term effectiveness of these systems in reducing the incidence of diarrhea. While boreholes have the potential to provide clean and safe water, their success in reducing waterborne diseases is often influenced by various factors, including maintenance, water quality, and community engagement. Many rural communities lack the technical expertise or financial resources to maintain borehole systems, leading to challenges such as contamination, pump failure, and water scarcity [8]. Furthermore, cultural and behavioral factors, such as inadequate sanitation practices and a lack of hygiene education, can undermine the potential benefits of borehole water systems. This study aims to explore the effectiveness of borehole water systems in reducing diarrhea in rural West Africa by reviewing existing evidence and identifying key factors that contribute to the success or failure of these interventions [9]. This study aims to evaluate the effectiveness of borehole water systems in reducing diarrhea incidence in rural West Africa and to explore the factors influencing their success and sustainability. It will examine the challenges faced by communities in maintaining these systems, the role of community engagement and hygiene education, and the overall health impact, particularly on vulnerable populations. One key objective is to assess how community involvement and hygiene education can enhance the effectiveness of borehole systems, ensuring that they are properly maintained and that communities adopt safe water practices. By analyzing the sustainability of these systems over time, the study will provide insights into long-term solutions for waterborne diseases. The research questions aim to identify the effectiveness of boreholes in reducing diarrhea, the factors influencing their success or failure, and the broader health impact of clean water access. This study is significant for guiding future water supply interventions in rural West Africa, as it highlights the importance of tailored, culturally appropriate solutions and integrated WASH (water, sanitation, and hygiene) interventions. The findings will also be valuable for policymakers, NGOs, and development practitioners, offering practical insights into improving public health outcomes in similar low-resource settings globally.

Borehole Water Systems: A Solution for Rural Water Supply

Boreholes, drilled deep into the ground to access groundwater, have become a common solution to the water scarcity and contamination challenges faced by rural communities in West Africa. These systems typically involve drilling wells into aquifers to provide clean, reliable water that is less likely to be contaminated by surface pathogens [10]. Borehole water systems are often considered more sustainable than surface water sources due to the relative purity of groundwater. Once installed, boreholes can provide year-round access to safe drinking water, reducing the reliance on surface water, which is more prone to contamination, especially during the rainy season. The success of borehole systems in rural areas is also attributed to their relatively low operational costs and their ability to provide clean water to a large number of households without the need for extensive infrastructure [11]. However, their effectiveness in reducing diarrheal diseases depends on several factors, including proper maintenance, water quality monitoring, and community involvement.

Effectiveness of Borehole Systems in Reducing Diarrhea

The effectiveness of borehole water systems in reducing the incidence of diarrhea in rural West Africa can be evaluated through multiple indicators, each reflecting the broader impact of access to clean water on public health. One of the primary measures of success is the reduction in diarrheal incidence. Numerous studies have demonstrated that communities with access to borehole water experience lower rates of diarrhea and other waterborne diseases [12]. For instance, research conducted in rural Ghana showed that the introduction of boreholes significantly reduced diarrhea cases among children under five, a group particularly vulnerable to waterborne illnesses. Similarly, studies from Sierra Leone revealed that borehole access resulted in lower rates of cholera and diarrhea compared to communities relying on unprotected water sources like rivers or open wells.

Improved water quality is another critical factor contributing to the effectiveness of borehole systems [13]. Groundwater, typically drawn from boreholes, is often free from contaminants such as bacteria, viruses, and fecal coliforms that are commonly found in surface water. Borehole water in rural West Africa frequently meets

international drinking water standards for microbial quality, making it a safer alternative to surface sources, which carry a higher risk of contamination and associated diseases [14].

The long-term health benefits of borehole systems are also noteworthy. With sustained access to clean water, communities experience a marked decline in waterborne diseases, fewer healthcare visits for diarrhea, and improved overall health outcomes. In some cases, there has been a significant reduction in child mortality rates, particularly in rural regions where diarrhea poses a substantial health threat [15]. Finally, borehole systems often lead to community empowerment and behavior change. By involving residents in the construction, management, and maintenance of water systems, these projects foster a sense of ownership and responsibility. This engagement encourages better hygiene practices, such as handwashing and safe water storage, which further mitigate the risk of diarrhea and other waterborne diseases [16].

Challenges and Limitations

Borehole water systems have proven to be an essential solution for improving water access in rural West Africa, but their effectiveness in reducing waterborne diseases like diarrhea is not without challenges. One of the primary limitations is the poor maintenance and sustainability of these systems [17]. Over time, boreholes can become non-functional due to mechanical breakdowns, lack of spare parts, or a shortage of technical expertise in the community. When regular maintenance is neglected, the borehole's water supply becomes unreliable, and contamination can occur, diminishing its health benefits. Another major issue is water quality. Although borehole water is typically cleaner than surface water, it is still vulnerable to contamination from both natural and human-made sources [18]. In some West African regions, borehole water can be tainted by high levels of nitrates from agricultural runoff, heavy metals, or naturally occurring arsenic. Without consistent monitoring and treatment, these contaminants pose significant health risks, potentially exacerbating issues beyond just diarrhea. Additionally, access to borehole water is not always equitable. Some rural communities, particularly remote or marginalized populations, may not have access to boreholes, leaving them reliant on unsafe water sources [19]. Even where boreholes are installed, their distance from communities can create barriers, as residents may have to travel long distances to collect water, which reduces the system's effectiveness. Finally, environmental and geographical factors play a critical role. In some areas, aquifers may be insufficient to meet the growing population's needs, especially during dry seasons. Over-exploitation of groundwater sources can lead to water shortages, further compromising the reliability and long-term sustainability of borehole systems [20]. These challenges underscore the need for comprehensive planning, investment in maintenance, and equitable distribution to ensure the effectiveness of borehole water systems in improving public health.

CONCLUSION

Borehole water systems have proven to be an invaluable resource for improving access to safe drinking water and significantly reducing the prevalence of diarrhea in rural West Africa. In areas where surface water is often contaminated with pathogens, boreholes offer a reliable alternative by providing clean and potable water, thus contributing to better public health outcomes. This access to safe water has led to a noticeable decline in waterborne diseases, particularly diarrhea, which remains a major health issue in many rural communities. However, for borehole water systems to continue delivering these benefits in the long term, several challenges must be addressed. These include the need for ongoing maintenance to prevent system breakdowns, ensuring consistent water quality through regular monitoring, and promoting equitable access for all community members, including marginalized populations. Furthermore, environmental sustainability must be factored into the management of these systems to ensure they are resilient to climate change and environmental degradation. With robust community involvement, effective monitoring programs, and a focus on sustainability, borehole systems can remain a critical tool in improving public health and reducing the burden of waterborne diseases like diarrhea in rural West Africa.

REFERENCES

1. Terefe, B., Jember, M.M., Assimamaw, N.T.: Access to drinking safe water and its associated factors among households in East Africa: a mixed effect analysis using 12 East African countries recent national health survey. *Journal of Health, Population and Nutrition*. 43, 72 (2024). <https://doi.org/10.1186/s41043-024-00562-y>
2. Catherine, A.N., Ayesiga, S., Rukundo, G.Z., Lejju, J.B., Byarugaba, F., Tamwesigire, I.K.: Community perceptions and practices on quality and safety of drinking water in Mbarara city, south western Uganda. <https://doi.org/10.1371/journal.pwat.0000075>
3. Alum, E. U., Obeagu, E. I., Ugwu, O. P. C. Curbing Diarrhea in Children below five years old: The sub-Saharan African Standpoint. *J. New Medical Innovations and Research*. 2024;5(1); DOI:10.31579/2767-7370/083

- 4.1 in 4 people globally still lack access to safe drinking water – WHO, UNICEF, <https://www.who.int/news/item/26-08-2025-1-in-4-people-globally-still-lack-access-to-safe-drinking-water---who--unicef>
5. Afonso, A.C., Saavedra, M.J., Gomes, I.B., Simões, M., Simões, L.C.: Current microbiological challenges in drinking water. *Journal of Water Process Engineering*. 72, 107614 (2025). <https://doi.org/10.1016/j.jwpe.2025.107614>
6. Uhama, K. C., Ugwu, O. P. C., Alum, E. U., (2024). Role of Phytochemical-Rich Foods in Mitigating Diarrhea among Diabetic Patients. *Research Invention Journal of Scientific and Experimental Sciences*. 3(1):45-55.
7. Omuna D., Obaroh I. O., Akiyode O. O., Eniru E. I., Tiyo C. E & Omoding, J. (2024). Impacts of climate change on water security in Uganda: A review. *Int. J. Adv. Multidiscip. Res.* 11(9): 47-60. DOI: <http://dx.doi.org/10.22192/ijamr.2024.11.09.005>
8. Agah, V. M., Orji, O. U., Ezeani, N. N., Ugwu, O. P., Bawa, I., Omang, W. A. and Itodo, M. O. (2023). Physico-chemical and Bacteriological Analysis of Water used for Drinking and other Domestic Purposes in Amaozara Ozizza, Afikpo North, Ebonyi State, Nigeria. *Nigerian Journal of Biochemistry and Molecular Biology*, 38(1), 1-8. <https://doi.org/10.2659/njbmb.2023.151>.
9. Ugwu, O. P. C., Alum, E. U., Uhama, K. C. (2024). Phytochemicals and Vitamins as Adjunct Therapies for Diarrhea in Diabetic Patients. *Research Invention Journal of Research in Medical Sciences*. 3(2):27-37.
10. Agyemang, V.O.: Groundwater resources development for a sustainable water supply in developing countries: A case study of Ghana. *Cleaner Water*. 4, 100104 (2025). <https://doi.org/10.1016/j.clwat.2025.100104>
11. Awor P, Wamani H, Bwire G, Jagoe G, Peterson S (2012). Private sector drug shops in integrated community case management of malaria, pneumonia, and diarrhea in children in Uganda. *The American journal of tropical medicine and hygiene*, 87, (5), 92, doi: [10.4269/ajtmh.2012.11-0791](https://doi.org/10.4269/ajtmh.2012.11-0791).
12. Cha, S., Kang, D., Tuffuor, B., Lee, G., Cho, J., Chung, J., et al: The Effect of Improved Water Supply on Diarrhea Prevalence of Children under Five in the Volta Region of Ghana: A Cluster-Randomized Controlled Trial. *International Journal of Environmental Research and Public Health*. 12, 12127–12143 (2015). <https://doi.org/10.3390/ijerph121012127>
13. Alum E U, Obeagu E I, Ugwu O P. C.. Enhancing quality water, good sanitation, and proper hygiene is the panacea to diarrhea control and the attainment of some related sustainable development goals: A review. *Medicine (Baltimore)*. 2024 Sep 20;103(38):e39578. doi: 10.1097/MD.00000000000039578. PMID: 39312342; PMCID: PMC11419503.
14. Mofokeng, R., Bezuidenhout, C.C., Van Wyk, D.A.B.: Toward inclusive microbial monitoring: The case for pathogenic yeast surveillance in South African groundwater. *Total Environment Microbiology*. 1, 100038 (2025). <https://doi.org/10.1016/j.temicr.2025.100038>
15. Critchley J. A, Ejemot-Nwadiaro R. I, Ehiri J. E, Arikpo D, Meremikwu M. M (2015). Hand washing promotion for preventing diarrhea. *Cochrane Database of Systematic Reviews*, 9, 9. Art. No.: CD004265. DOI: 10.1002.
16. Ogban G. I, Ndueso E. M, Iwuafor A. A, Emanghe U. E, Ushie S. N, Ejemot-Nwadiaro R. I (2020). Basic Knowledge of Childhood Diarrhea and Health-seeking Practices of Caregivers of Under-five Children in Calabar-South, Calabar, Nigeria. *Asian Journal of Medicine and Health*, 18, (4), 12-23. <https://doi.org/10.9734/ajmah/2020/v18i430195>.
17. Chimombo, M.J.: Frontiers | Bridging the divide: analysing disparities in drinking water services between urban and rural areas in Lilongwe, Malawi—implications for water justice and infrastructure equity. <https://doi.org/10.3389/frwa.2025.1615202>
18. Ogban G. I, Ndueso E. M, Iwuafor A. A, Emanghe U. E, Ushie S. N, Ejemot-Nwadiaro R. I (2021). Availability and Affordability of Sanitation and Health Promoting Amenities: Driving the Impact of Knowledge of Childhood Diarrhea on Health-Seeking Practices of Under-five ... *Highlights on Medicine and Medical Research*, 3. B P International, 140-155. <https://doi.org/10.9734/bpi/hmmr/v3/2344E>.
19. Nlemolisa, O.R., Ogbulie, J.N., Orji, J.C., Nweke, C.O., Kemka, U.N., Gaius-Mbalisi, V.K., Ihenetu, F.C.: Groundwater contamination and health risks near waste dumps and mechanic workshops: A seasonal perspective. *Cleaner Water*. 4, 100090 (2025). <https://doi.org/10.1016/j.clwat.2025.100090>
20. Grönwall, J., Danert, K.: Regarding Groundwater and Drinking Water Access through A Human Rights Lens: Self-Supply as A Norm. *Water*. 12, 419 (2020). <https://doi.org/10.3390/w12020419>

CITE AS: Mugisha Emmanuel K. (2026). The Effectiveness of Borehole Water Systems in Reducing Diarrhea in Rural West Africa: A Review. NEWPORT INTERNATIONAL JOURNAL OF PUBLIC HEALTH AND PHARMACY, 7(2):14-18. <https://doi.org/10.59298/NIJPP/2026/721418>