**©IDOSR PUBLICATIONS** 

International Digital Organization for Scientific Research IDOSR JOURNAL OF SCIENTIFIC RESEARCH 10(1):62-70, 2025. https://doi.org/10.59298/IDOSRJSR/2024/10.1.62700

## Omagor and Abdi ISSN: 2550-794X

IDOSRJSR10.1.62700

## Prevalence and Factors Associated with Severe Malaria in Children Under 5 Years of Age at Hoima Regional Referral Hospital, Western Uganda

## **Omagor Herbert and Said Abdi**

Faculty of Clinical Medicine and Dentistry Kampala International University, Uganda.

#### ABSTRACT

This study aimed to determine the prevalence and factors associated with severe malaria in children under five years of age at Hoima Regional Referral Hospital (HRRH) in western Uganda. A cross-sectional descriptive study was conducted in the pediatric ward of HRRH among 288 caregivers of children under five years admitted to the ward. Data were collected using a structured questionnaire, coded, entered, and analyzed using SPSS version 25. The majority of the children were female and above two years of age. Most caregivers were farmers, and many had no formal education. Among the 288 children admitted to the pediatric ward, 188 (65.3%) had severe malaria. Factors significantly associated with severe malaria included age and distance from home to the hospital. Children older than two years had a significantly higher likelihood of severe malaria (P  $\leq$  0.05), with the odds being twice as high compared to those aged two years or younger. Additionally, children living five kilometers or more from the hospital had twice the odds of having severe malaria compared to those living within one kilometer. The odds increased fourfold for children residing between two and five kilometers from the hospital. The prevalence of severe malaria was high among children admitted to HRRH. Significant risk factors included being older than two years and living more than two kilometers from the health facility.

Keywords: Severe malaria, Children under five, Prevalence, Risk factors, Hoima Regional Referral Hospital

## INTRODUCTION

Malaria is one of the most common and lifethreatening tropical diseases worldwide  $\lceil 1 \rceil$ . It is caused by Plasmodium parasites, which are transmitted through the bite of an infected female Anopheles mosquito, primarily between evening and morning [2]. In humans, malaria is mainly caused by four species of Plasmodium: Plasmodium falciparum, vivax, Plasmodium Plasmodium malariae, and Plasmodium ovale. Among these, Plasmodium falciparum and Plasmodium vivax are the most prevalent [3,4]. The majority of malaria cases are concentrated in sub-Saharan Africa, where Plasmodium falciparum accounts for nearly all infections. It is the most virulent species, responsible for over 90% of malaria-related deaths worldwide, making it a significant public health concern [5]. Globally, malaria remains widespread in tropical regions, with approximately 3.4 billion people at risk each year, including 1.2 billion at high risk [6]. Despite being preventable and treatable, malaria continues to cause significant morbidity and mortality, particularly in resource-limited settings.

An estimated 300-500 million people suffer from malaria annually, resulting in 1.5-2.7 million deaths, with sub-Saharan Africa contributing to over 80% of the global malaria mortality burden. Children under five years and pregnant women are the most vulnerable groups. Malaria is the leading cause of death in children under five, accounting for 77% of all malaria-related deaths globally [7]. In Africa, a child dies from malaria every minute. Severe malaria often results from delayed treatment and poor management of uncomplicated malaria  $\lceil 8 \rceil$ . It is defined as the presence of clinical and laboratory evidence of vital organ dysfunction. Nearly all malaria deaths are due to Plasmodium falciparum infection [9]. Although new drugs and vaccines have been introduced, malaria remains a major health challenge, and current strategies focus on prevention and control [10]. Uganda has one of the highest malaria burdens in sub-Saharan Africa, ranking third in Plasmodium falciparum infections and among the highest in malaria transmission rates globally [11]. Malaria transmission is stable and perennial in 90-

95% of the country. According to Uganda's Health Management Information System (HMIS), malaria accounted for 34% of outpatient visits and 28% of hospital admissions in 2015. Of these reported cases, 55% were laboratory-confirmed, increasing to 64% between July and December, showing an improvement in case management [12]. Efforts to reduce malaria transmission have been enhanced through interventions such as intermittent preventive treatment for pregnant women, distribution of long-lasting insecticide-treated nets (LLINs), and early diagnosis and treatment [4]. Despite these efforts, malaria remains the leading cause of death in Uganda, accounting for over 27% of all fatalities [13]. Uganda also has the highest malaria incidence worldwide, with 478 cases per 1,000 people annually [14]. In 2016, there were an estimated 216 million malaria cases globally, leading to 445,000 deaths, with 91% of these occurring in sub-Saharan Africa [15]. Children under five accounted for two-thirds of malaria-related deaths that year. According to Uganda's Ministry of Health, malaria accounts for 30-50% of outpatient visits, 15-

#### **Study Design**

This was a cross-sectional study in which quantitative data was collected.

## **Study Setting**

The study was conducted in the pediatric ward of Hoima Regional Referral Hospital (HRRH), located approximately 180 km north of Mbarara, the largest city in the western region of Uganda. Hoima is situated about 297 km west of Kampala, Uganda's capital city, along a fully tarmacked two-lane highway. Hoima serves as the main town of the Bunyoro Kingdom in Hoima District. The district is bordered by Masindi and Buliisa Districts to the north, Kyankwanzi District to the east, and Kibaale District to the south, extending to the national boundary with the Democratic Republic of Congo in the west. HRRH is a public hospital funded by the Uganda Ministry of Health, and general healthcare services at the hospital are provided free of charge.

## **Study Population**

The study population consisted of children under five years of age who were admitted to the pediatric ward of Hoima Regional Referral Hospital.

#### **Inclusion Criteria**

Children below five years of age admitted to the pediatric ward of Hoima Regional Referral Hospital. Caregivers who consented to participate in the study.

## **Exclusion Criteria**

Children below five years of age admitted to the pediatric ward but who died on admission.

Children admitted with severe malaria but older than five years of age.

#### Omagor and Abdi

20% of hospital admissions, and 19-14% of inpatient mortality [16]. Malaria-related mortality in children under five often exceeds 10%, increasing with disease severity. Severe malaria risk factors include coma, convulsions, acidosis, respiratory distress, hyperpyrexia, elevated lactate hypoglycemia, concentrations, and high levels of tumor necrosis factor [17]. Although substantial efforts have been made by the international community, Uganda's Ministry of Health, and healthcare services to expand malaria prevention, diagnostics, and treatment, severe malaria remains a significant public health problem. Many children under five continue to suffer from severe malaria, particularly those admitted to Hoima Regional Referral Hospital (HRRH). However, no previous studies have been conducted in the pediatric ward of HRRH to identify the risk factors associated with severe malaria. Therefore, this study aims to determine the prevalence and factors associated with severe malaria in children under five years at Hoima Regional Referral Hospital in the Bunyoro region.

## METHODOLOGY

## Sample Size Determination

The sample size was determined using the Kish-Leslie formula:

 $n = z^2 p (1-p) / E^2$ Where:

- n = Estimated minimum sample size required

- P = 75% prevalence of severe malaria in children

under five years in sub-Saharan Africa [15]

- Z = 1.96 (for a 95% Confidence Interval)
- E = Margin of error set at 5%

 $n = 1.96^2 \times 0.75(1 - 0.75) / 0.05^2$ 

n = 288

#### Sampling Procedure

All children under five years of age who met the inclusion criteria were enrolled until the target sample size was reached.

#### **Data Collection Procedure**

Structured questionnaires were used to interview caregivers of children with severe malaria. The majority of the questions were a mix of open-ended and closed-ended formats. Data was collected on both caregivers and children with severe malaria. The questionnaire was designed to capture demographic information on caregivers and their children, along with healthcare system and household factors. Caregivers of children with severe malaria were interviewed on the day of admission or the next day, once the child's health had stabilized. Trained research assistants administered the questionnaires.

#### Data Analysis

Following data collection, the data was entered into SPSS version 20 for analysis. Each independent

variable was analyzed in a univariate form using tabular presentation. Subsequently, independent variables were compared with the dependent variable in a bivariate analysis. Finally, multivariate analysis was conducted to establish conclusions for this research.

## **Quality Control Measures**

Study procedures were taught to selected research assistants. The accuracy, consistency, and completeness of completed questionnaires were evaluated on a daily basis. Pretesting of the questionnaires was also conducted to ensure reliability.

## **Ethical Considerations**

Ethical approval was obtained from KIU IREC and HRRH. The study proposal was reviewed and approved by the HRRH Department of Pediatrics before commencement.

## Omagor and Abdi

#### Confidentiality

The confidentiality of participants' information was ensured through the use of unique reference codes during data collection and analysis. Participants were assured of complete confidentiality regarding the information they provided.

#### **Risks and Benefits**

There were no painful procedures involved, except for treatment purposes. Children received treatment before participation in the research.

#### **Limitations and Delimitations**

1. The study was prone to selection bias as it was conducted at a health facility.

2. Non-cooperative respondents limited the scope of the study, and recall bias may have affected data accuracy.

3. The diversity of languages spoken by patients admitted to Hoima Regional Referral Hospital may have affected the accuracy of data collection.

4. Referral bias may have influenced the findings, as most children admitted to a referral hospital are likely to have severe malaria.

Variable	Frequency (n)	Percentage (%)
Age of the child		
Below 2 years	112	38.9
Above 2 years	176	61.1
Sex of a child		
Male	135	46.9
Female	153	53.1
Occupation of the caregiver		
Peasant farmer	201	69.8
Not a farmer	87	30.2
Education level of the caregiver		
None	127	44.1
Primary	107	37.2
Secondary	36	12.5
Tertiary	18	6.2

RESULTS Socio-demographic findings

#### Prevalence of severe malaria among children under five years at HRRH Table 2: Malaria status of children admitted at HRRH

Malaria diagnosis	Frequency (n)	Percentage (%)
Severe malaria Not severe un complicated	188	65.3
malaria	100	34.7

Table 2 above shows that out of the 288 children admitted on pediatric ward at FRRH in the study, 188 (65.3%) had severe malaria.

Variable	Severe n (%)	ncomplicated	adjusted OR (95%	P-value
		(%)	CI)	
Age of the child				
Above 2 years	79(70.5)	33(29.5)	2.1 (1.15-3.93)	0.015
Below 2 years	111(63.1)	67(26.9)	1.0	
Sex of a child	· · ·			
Male	115(85.2)	20(14.8)	1.0	
Female	73(47.7)	80 (52.3)	3.8(0.82-17.54)	0.087
Occupation of the caregiver				
Peasant farmer	116(57.7)	84(42.3)	1.0	
Not a farmer	71 (81.6)	16(18.4)	1.0(0.49-2.08)	0.976
Education level of the caregiver	× ,		· · · · ·	
None	81(63.8)	46(36.2)		
Primary	61(57.0)	46(43.0)	3.00(0.62 - 14.45)	0.172
Secondary	31(86.1)	5(13.9)	2.67(0.57-12.47)	0.214
Tertiary	15(83.3)	3(16.7)	2.68(0.53-13.54)	0.232

r D: • 1.: **f**\_\_\_\_\_ т D ſ C J

From the table 3 above only child's age and sex age had P-values <0.2 thus taken for the multivariate analysis.

Multivariate binary logistic regression of socio demographic factors associated with severe malaria among the study participants.

Table 4: Multivariate binary logistic regression of socio demographic factors associated with severe malaria among the study participants.

Variable	Adjusted OR (95% CI)	P-value
<b>Age of the child</b> Above 2 years Below 2 years	2.0 (1.07-3.74) 1.0	0.030
<b>Sex of a child</b> Female Male	1.84(0.97 <b>-</b> 3.50) 1.0	0.063

Following multivariate analysis, being a child above 2 years was significantly associated with severe malaria (P-value  $\leq 0.05$ ). The odds of having severe malaria were two-fold higher among children aged above 2 years as compared to those below 2 years.

# Household factors associated with severe malaria in children under five years admitted on pediatric ward at HRRH

## Table 5: Results of Bivariate Analysis for household factors

Variable severe Not Un adjusted OR P-value n (%) severe (95% CI) n (%)

Household size				
Below 5 in number	80(80.0)	20(20.0)	3.4(0.67 - 17.09)	0.142
Above 5 in number	108(57.4)	80(42.6)	1.0	
Do you have mosquito nets at home				
Yes	37(32.5)	77(67.5)	1.0	
No	121(78.6)	33(22.2)	1.5(0.78 - 2.78)	0.233
Number of under five children at home				
One child	87(56.1)	55 (43.9)	1.0	
2 or more children	101(69.2)	45 (30.1)	1.7(0.72-4.05)	0.219
Type of family				
Polygamous	143 (88.6)	21(14.4)	1.00	
Monogamous	45 (36.3)	79(63.7)	3.6 (0.82-15.89)	0.091
Materials used to wall of your house				
Wattle and mud Wood	97~(68.8)	44(31.2)	1.00	
	43(70.5)	18(29.5)	1.1 (0.39-3.25)	0.814
Sand and bricks	48(55.8)	38(44.2)	1.5(0.47- $4.91)$	0.490

In the Bivariate analysis in table 5 above household size and type of family had P-values <0.2 thus taken for the multivariate analysis.

## Multivariate binary logistic regression: Household factors associated with severe malaria among the study participants.

From table 6 below, both household size and type of family were not statistically significant in relation to severe malaria since both P-values > 0.05.

## Omagor and Abdi

## www.idosr.org

Table 6: Model one of multivariate binary logistic regression: Household factors associated with severe malaria among the study participants

Variable	Adjusted OR (95% CI)	P-value
Household size		
Below 5 in number	1.80(0.56-5.76)	0.320
Above 5 in number	1.00	
Type of family		
Polygamous	1.00	
Monogamous	3.00(0.62 - 14.45)	0.172

## Health related factors associated with severe malaria in children under five years admitted on pediatric ward at HRRH.

From the table 7 below distance from home to the health facility and waiting time before taking a child to hospital had P-values <0.2 thus taken for the multivariate analysis.

Severe Variable	Not severe	Unadjusted OR P-value	n (%)	(95% CI)
Community health facility yes no		II (70)		
Community hearth facility yes no	109(709)	07 (00 0)	1.0	
	103(79.2)	27(20.8)	1.0	0.079
Distance from home to the health facility	89(93.4)	73 (40.0)	1.0 (0.30-2.85)	0.973
9-5 kilometers				
1 kilometer	60 (49 8)	63(519)	4.0(1.64-9.83)	0.00%
> 5 kilometers	93(76.9)	98(931)	1.0	0.002
	35(91.3)	9(8.7)	2.0(1.02 - 3.93)	0.046
Means of transport to hospital	00 (0110)	0 (0.1)	210 (1102 0100)	01010
Foot	66(54.1)	56(45.9)	1.0	
Bicvcle	11(40.7)	16(59.3)	1.4(0.29-6.88)	0.664
Boda boda	90(81.8)	20(18.2)	2.2(0.201 - 1.170)	0.207
Motor car	21(72.4)	8(27.6)	0.8 (0.278-2.409)	0.718
Medication given to child when identifies	( <i>)</i>	( )	( /	
sickness				
no medication given	101(65.2)	54(34.8)	1.00	
local herbs	37(52.1)	34(47.9)	0.86 (0.32-2.34)	0.758
medication from drug shops/pharmacy	50(80.6)	12(19.4)	0.84(0.29-2.42)	0.743
Waiting time before taking a child to hospital				
Within one day	79(81.4)	18(18.6)	1.6(0.126 - 1.00)	0.150
After one day	109(57.1)	82(42.9)	1.0	0.100
Time taken to see a health worker at the			-	
hospital				
with 1 hour	53(71.3)	27(29.7)	0.63(0.18 - 2.24)	0.472
with 2 hours	77(67.0)	38(33.0)	0.74 (0.45-1.24)	0.261
> 3 hours	58(62.4)	35(37.6)	1.00	
Getting all medication when at health facility	× /	× /		
Within one day	60(84.5)	11(15.5)	0.93(0.53 - 1.62)	0.790
After one day	128(59.0)	89(31.0)	1.00	

## Multivariate binary logistic regression: Health related factors associated with severe malaria among the study participants

According to table 8, distance of 2-5 kilometers from home to the hospital and distance of greater than 5 kilometers from home to the hospital were significantly associated with severe malaria. The odds of having severe malaria were two-fold higher among children within 5 kilometers or greater from

#### Omagor and Abdi

home to the hospital as compared to distance of 1 kilometer from home to the hospital. In addition, the odds of having severe malaria were four-fold higher among children taking a distance of 2-5 kilometers from home to the hospital as compared to those taking distance of 1 kilometer from home to the hospital.

Table 8: Final model of multivariate binary logi	stic regression	: Health related	l factors a	associated	with
severe malaria among the study participants.					

Variable	Adjusted OR (95% CI)	P-value
e from home to the health facility		
2-5 kilometers	3.8 (1.51-9.56)	0.005
1 kilometer	1.0	
> 5 kilometers	2.1 (1.06-4.27)	0.033
<b>g time before taking a child to hospital</b> Within one day After one day	1.1(0.37 <b>-</b> 2.98) 1.00	0.924

#### DISCUSSION

### Prevalence of Severe Malaria in Children Under Five Years of Age

This study found that the prevalence of severe malaria among children admitted to the pediatric ward at FRRH was 65.3%. In contrast, a study by Zgambo et al., [18] in Malawi reported an increase in the prevalence of severe malaria among children under five from 28% in 2012 to 33% in 2014. The prevalence observed in this study is significantly higher. According to UBOS, malaria contributes to 20-30% of all pediatric hospital admissions nationwide, and this study indicates that 65.3% of these cases are severe [19]. Consequently, if this trend continues, the national under-five malaria mortality rate may increase, potentially hindering Uganda's progress toward achieving the Sustainable Development Goals (SDGs).

## Association Between Socio-Demographic Factors and Severe Malaria in Children Under Five Years of Age

This study identified a significant association between a child's age and the likelihood of severe malaria. Children older than two years had twice the odds of developing severe malaria compared to those under two years. This finding aligns with studies conducted in Uganda [20] and Malawi [18], both of which reported an increased risk of severe malaria among older children. This trend may be attributed to older children having more freedom to move outdoors at night, increasing their exposure to mosquito bites [21-26]. Additionally, Zgambo et al., [18] suggested that younger children are often covered well under mosquito nets as they share beds with their mothers, thereby reducing their exposure. However, this study's findings contrast with a study conducted in Rwanda by Nyirakanani et al., [22], which reported a higher prevalence of severe malaria among children aged 1 to 12 months compared to those aged 13 to 59 months. This discrepancy could be due to differences in the age groups considered in each study [24-27].

While various studies have demonstrated a significant association between household factors and severe malaria-such as household size, number of children under five years, family structure, electricity availability, and access to treated mosquito nets-this study found no significant association between household factors and severe malaria [24-28].

## Association Between Health Facility Factors and Severe Malaria in Children Under Five Years of Age

This study identified a significant association between the distance to a health facility and severe malaria. Children residing five or more kilometers from a hospital had twice the odds of developing severe malaria compared to those living within one kilometer. Furthermore, children who lived 2-5 kilometers away had four times the odds of developing severe malaria compared to those within one kilometer of a hospital. These findings are consistent with a study in the Asikuma-Odoben-Brakwa District, Ghana, which reported that longer distances to health facilities encourage reliance on traditional medicine, allowing mild malaria cases to progress to severe forms [23]. The strong association between distance and severe malaria

malaria.

underscores the role of healthcare accessibility in malaria outcomes. Caregivers' treatment choices are often influenced by the availability and accessibility

In conclusion, this study highlights a high prevalence

of severe malaria among children admitted to the

pediatric ward at FRRH. The key factors

significantly associated with severe malaria were a

child's age and the distance from home to a

healthcare facility. Older children (above two years)

and those residing more than two kilometers from a

hospital were at a higher risk of developing severe

Inty CONCLUSION

#### Recommendations

of healthcare facilities, making longer distances a

critical risk factor for severe malaria.

Mothers and caregivers should be particularly vigilant about the health of children older than two years, as they are at increased risk of severe malaria.
The government and healthcare policymakers should prioritize establishing health centers closer to communities to improve accessibility.

- Academicians, researchers, and health organizations should conduct further studies to explore additional factors influencing severe malaria in children under five years of age.

#### REFERENCES

- Kungu, E., Inyangat, R., Ugwu, O. P. C., Alum, E. U. (2023). Exploration of Medicinal Plants Used in the Management of Malaria in Uganda. Newport International Journal of Research in Medical Sciences. 4(1):101-108.
- Alum EU, Tufail T, Agu PC, Akinloye DI, Obaroh IO. (2024). Malaria pervasiveness in Sub-Saharan Africa: Overcoming the scuffle. Medicine. 103(49), e40241. doi: 10.1097/MD.000000000040241. PMID: 39654176.
- Egwu CO, Aloke C, Chukwu J, Agwu A, Alum E, Tsamesidis I, et al. (2022). A world free of malaria: It is time for Africa to actively champion and take leadership of elimination and eradication strategies. Afr Health Sci., 22(4):627-640. doi: 10.4314/ahs.v22i4.68.
- 4. Ekpono EU, Aja PM, Ibiam UA, Alum EU, Ekpono UE. Ethanol Root-extract of Sphenocentrum jollyanum Restored Altered Haematological Markers Plasmodium in berghei-infected Mice. Earthline Journal of Chemical Sciences. 2019: 2(2): 189-203. https://doi.org/10.34198/ejcs.2219.189203.
- Alum EU, Ugwu OPC, Egba SI, Uti DE, Alum BN. (2024). Climate Variability and Malaria Transmission: Unraveling the Complex Relationship. INOSR Scientific Research. 11(2):16-22. https://doi.org/10.59298/INOSRSR/2024/1.1. 21622
- World Health Organization. (2015). World Malaria Report 2015. World Health. https://doi.org/ISBN 978 92 4 1564403
- Sarfo JO, Amoadu M, Kordorwu PY, Adams AK, Gyan TB, Osman AG, Asiedu I, Ansah EW. (2023). Malaria amongst children under five in sub-Saharan Africa: a scoping review of prevalence, risk factors and preventive interventions. Eur J Med Res., 28(1):80. doi: 10.1186/s40001-023-01046-1. PMID: 36800986; PMCID: PMC9936673.

- Mousa A, Al-Taiar A, Anstey NM, Badaut C, Barber 8. BE, Bassat Q, Challenger JD, Cunnington AJ, Datta D, Drakeley C, Ghani AC, Gordeuk VR, Grigg MJ, Hugo P, John CC, Mayor A, Migot-Nabias F, Opoka RO, Pasvol G, Rees C, Reyburn H, Riley EM, Shah BN, Sitoe A, Sutherland CJ, Thuma PE, Unger SA, Viwami F, Walther M, Whitty CJM, William T, Okell LC. (2020). The impact of delayed treatment of uncomplicated P. falciparum malaria on progression to severe malaria: A systematic review and a pooled multicentre individual-patient meta-analysis. PLoS Med. 17(10):e1003359. doi: 10.1371/journal.pmed.1003359. PMID: 33075101; PMCID: PMC7571702.
- 9. World Health Organization (WHO). (2012). Management of severe malaria.
- El-Moamly, A.A., El-Sweify, M.A. (2023). Malaria vaccines: the 60-year journey of hope and final success—lessons learned and future prospects. *Trop Med Health* 51, 29. https://doi.org/10.1186/s41182-023-00516-w
- Ranjbar, M., Tegegn Woldemariam, Y. (2024). Nonfalciparum malaria infections in Uganda, does it matter? A review of the published literature. *Malar* J 23, 207. https://doi.org/10.1186/s12936-024-05023-9
- Usaid. (2013). President's Malaria Initiative Uganda Malaria Operational Plan FY 1-45. Retrieved from http://www.pmi.gov/countries/mops/fy13/liberia \_mop\_fy13.pdf
- Katushabe, J., Nnyanzi, J.B. & Muwanga, G.S. (2024). Exploring the role of spending on malaria incidence in Uganda using the auto-regressive distributed lag approach. *Malar J* 23, 129. https://doi.org/10.1186/s12936-024-04929-8
- Zalwango, M.G., Zalwango, J.F., Kadobera, D. et al. (2024). Evaluation of malaria outbreak detection methods, Uganda, 2022. Malar J 23, 18. https://doi.org/10.1186/s12936-024-04838-w

#### Omagor and Abdi

- World Health Organization. (2015). World Malaria Report 2015. World Health. https://doi.org/ISBN 978 92 4 1564403
- Ministry of Health, (2014). Ministry of Health, Republic of Uganda. National Malaria Control Programme.
- von Seidlein, L., Olaosebikan, R., Hendriksen, I. C. E., Lee, S. J., Adedoyin, O. T., Agbenyega, T., Dondorp, A. M. (2012). Predicting the Clinical Outcome of Severe Falciparum Malaria in African Children: Findings from a Large Randomized Trial. *Clinical Infectious Diseases*, 54(8), 1080–1090. https://doi.org/10.1093/cid/cis034
- Zgambo, M., Mbakaya, B. C., & Kalembo, F. W. (2017). Prevalence and factors associated with malaria parasitaemia in children under the age of five years in Malawi: A comparison study of the 2012 and 2014 Malaria Indicator Surveys (MISs). *PLoS ONE*, *12*(4), 1–13.

https://doi.org/10.1371/journal.pone.0175537

- Uganda Ministry of Health. (2014). The Uganda Malaria Reduction Strategic Plan. 1–83.
- Roberts, D., & Matthews, G. (2016b). Risk factors of malaria in children under the age of five years old in Uganda. Malaria Journal. https://doi.org/10.1186/s12936-016-1290-x 45
- Moshi, I.R., Manderson, L., Ngowo, H.S. et al. (2018). Outdoor malaria transmission risks and social life: a qualitative study in South-Eastern Tanzania. Malar J 17, 397. https://doi.org/10.1186/s12936-018-2550-8
- 22. Nyirakanani, C., Chibvongodze, R., Habtu, M., Masika, M., Mukoko, D., & Njunwa, K. J. (2018). Prevalence and risk factors of asymptomatic malaria among under-five children in Huye District,

Southern Rwanda. Tanzania Journal of Health Research, 20(1), 1-7. https://doi.org/10.4314/thrb.v20i1.6

 Amegbor, P. M. (2017). An Assessment of Care-Seeking Behavior in Asikuma-Odoben-Brakwa District: A Triple Pluralistic Health Sector. Approach.

https://doi.org/10.1177/2158244017710688

- 24. Ugo Kungu Erisa, Inyangat Raphael, Ugwu Okechukwu Paul-Chima and Alum Esther (2023). Exploration of Medicinal Plants Used in the Management of Malaria in Uganda. NEWPORT INTERNATIONAL JOURNAL OF RESEARCH IN MEDICAL SCIENCES, 4(1): 101-108.
- Obeagu EI, EU Alum, OPC Ugwu (2023). The Gatekeeper of Iron in Malaria Resistance. Newport International Journal of Research in Medical Sciences, 4(2): 1-8.
- Yusuf S. Enechi, O.C., Ugwu, Kenneth K., Ugwu Okechukwu P.C. and Omeh (2013). Evaluation of the antinutrient levels of *Ceiba pentandra* leaves. 3(3): 394-400.
- 27. Esther U Alum, Okechukwu PC Ugwu, Patrick M Aja, Emmanuel I Obeagu, Joseph E Inya, Amarachi P Onyeije, Esther Agu, Chinaza Godswill Awuchi (2023). Restorative effects of ethanolic leaf extract of *Datura stramonium* against methotrexate-induced hematological impairments. *Cogent Food & Agriculture*, 9(1): 2258774.
- Emmanuel Ifeanyi Obeagu, Yahye Abdi Ahmed, Getrude Uzoma Obeagu, UO Bunu, OPC Ugwu, Esther U Alum (2023). Biomarkers of breast cancer: Overview. 8(1): 8-16.

CITE AS: Omagor Herbert and Said Abdi (2025). Prevalence and Factors Associated with Severe Malaria Children Under 5 Years of Age at Hoima Regional Referral Hospital, Western Uganda. IDOSR JOURNAL ( SCIENTIFIC RESEARCH 10(1):62-70. <u>https://doi.org/10.59298/IDOSRJSR/2024/10.1.62700</u>

Omagor and Abdi