

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/378012808>

EVALUATION OF LIPID PROFILE OF PATIENTS WITH DIABETES ATTENDING ABIA STATE UNIVERSITY TEACHING HOSPITAL ABA, NIGERIA

Article · February 2024

CITATIONS

0

READS

17

13 authors, including:



Harrison U. Nwanjo

80 PUBLICATIONS 881 CITATIONS

SEE PROFILE



Augustine U Obi

Imo State University

14 PUBLICATIONS 67 CITATIONS

SEE PROFILE



Emmanuel Ifeanyi Obeagu

Kampala International University (KIU)

1,329 PUBLICATIONS 8,468 CITATIONS

SEE PROFILE



EVALUATION OF LIPID PROFILE OF PATIENTS WITH DIABETES ATTENDING ABIA STATE UNIVERSITY TEACHING HOSPITAL ABA, NIGERIA

C. O. Ahiara^{*1}, I. F. Onyeakplam¹, D. C. Nwosu¹, H. U. Nwanjo¹, C. O. Chima¹, G. C. Anyanwu¹, F. C. Emegaha¹, O. C. Okamgba², J. O. Kabiri¹, I. O. Onuh¹, I. C. Amobi¹, A. U. Obi³ and Emmanuel Ifeanyi Obeagu^{*4}

¹Department of Medical Laboratory Science, Imo State University, Owerri, Imo State, Nigeria.

²Department of Medical Laboratory Science, Abia State University Uturu, Abia State, Nigeria

³Department of Anatomy and Neurobiology, Imo State University Owerri, Imo State, Nigeria.

⁴Department of Medical Laboratory Science, Kampala International University, Uganda.



***Corresponding Author: C. O. Ahiara**

Department of Medical Laboratory Science, Imo State University, Owerri, Imo State, Nigeria.

Article Received on 21/12/2023

Article Revised on 11/01/2024

Article Accepted on 01/02/2024

ABSTRACT

The evaluation of lipid profile in diabetic patients attending Abia State Teaching Hospital Aba, was carried out. Venous blood samples were collected from 150 participants who gave consent. This comprises 100 diabetic patients as test and 50 healthy subjects as control. Cholesterol, Triglyceride, High density lipoprotein and Low-density lipoprotein were determined using semi-auto analyzer (Rato 9200). Data from this study were analyzed using statistical package for the social sciences (SPSS). Result shows that, serum cholesterol (4.96 ± 0.91 mmol/l), triglyceride (1.84 ± 0.42 mmol/l), low density lipoprotein (2.40 ± 0.66 mmol/l) levels were higher in study population compared with control (4.48 ± 0.41 mmol/l, 1.63 ± 0.09 mmol/l, 1.67 ± 0.22 mmol/l, $176.18 \pm 25.26 \times 10^9$ /l, 11.85 ± 0.63 secs, respectively) ($p < 0.05$ in each case). From the findings, management of conditions related to cardiovascular disease, arteriosclerotic disease, anemia and stress in diabetics may benefit patients if lipid profile, and some hematological parameters are included as part of their routine laboratory investigations.

KEYWORDS: Lipid Profile, Diabetes mellitus, cardiovascular disease, arteriosclerotic disease.

INTRODUCTION

Diabetes mellitus (DM), is a group of metabolic disorders in which there is high blood sugar level over a prolonged period and it is commonly referred to as diabetes.^[1-6] Frequent urination, increased thirst, and increased hunger are symptoms of high blood sugar.^[7-12] Many complications are resulted as a cause of untreated diabetes.^[13-18] Diabetic ketoacidosis, hyperosmolar hyperglycemic state, or death are as a result of acute complications. However, cardiovascular disease, stroke, chronic kidney disease, foot ulcers, and damage to the eyes are included as long-term complication.^[19-24]

Diabetes mellitus results from either the pancreas is not generating enough insulin or the cells of the body do not respond appropriately to the insulin produced.^[19]

Reduced HDL cholesterol, a predominance of small dense LDL particles, and elevated triglyceride levels are abnormalities associated with a clustering of interrelated plasma lipid and lipoprotein in Insulin resistance and form 2 diabetes.^[1] Every single of these dyslipidemic

structures is related with an amplified risk of cardiovascular disease. Increased hepatic secretion of large triglyceride-rich VLDL and impaired clearance of VLDL appears to be of central importance in the pathophysiology of this dyslipidemia. Small dense LDL particles arise from the intravascular processing of specific larger VLDL precursors. Typically, reduced plasma HDL levels in type 2 diabetes are manifest as reductions in the HDL2b subspecies and relative or absolute increases in smaller denser HDL3b and HDL3c. Although behavioral interventions such as diet and exercise can improve diabetic dyslipidemia, for most patients, pharmacological therapy is needed to reach treatment goals.

All the abnormalities occur in many patients despite normal LDL cholesterol levels. These changes are also a feature of the insulin resistance syndrome (also known as the metabolic syndrome), which underlies many cases of type 2 diabetes.^[1] In fact, pre-diabetic individuals often exhibit an atherogenic pattern of risk factors that includes higher levels of total cholesterol, LDL

cholesterol, and triglycerides and lower levels of HDL cholesterol than individuals who do not develop diabetes. Insulin resistance has striking effects on lipoprotein size and subclass particle concentrations for VLDL, LDL, and HDL.

MATERIALS AND METHODS

STUDY AREA

The study was carried out at Abia State University Teaching Hospital (ABSUTH), Aba city in Abia state, South East of Nigeria.

STUDY POPULATION

The size of population was calculated using the method of Aroye 2004 with the formula $n = (z^2 pq / d^2)$, and one hundred and fifty(150) subjects were recruited into the study. This comprises fifty(50) non-diabetic subjects as control and one hundred(100) diabetic subjects as test subjects.

SELECTION CRITERIA

INCLUSION; Those selected are;

- (i) Male and Female subjects of age 18 years to 74 years,
- (ii) Diabetic patients with blood sugar 10mmol/l and above,
- (iii) The subjects that gave their consent.

EXCLUSION; The excluded subjects are;

- (i) Male and Female subjects below the age of 18 years
- (ii) Subjects of blood sugar below 10mmol/l
- (iii) The subjects that did not give consent.

LABORATORY PROCEDURES

The reagents were commercially purchased and the manufacturers' standard operating procedures (S.O.P) were strictly adhered to.

DETERMINATION OF CHOLESTEROL: (enzymatic endpoint method) of Turfitt 1994.

Assay procedure; 1000µl of cholesterol liquor and 10µl of sample, standard and control were added into test tubes respectively, and incubated for 10 minutes at 37°C. The absorbance of standard and test were read (measured)

against reagent blank within 60 minutes at 500nm wavelength.

$$\text{Conc. of cholesterol} = \frac{\text{Abs. of test} \times \text{Conc. of Std.}}{\text{Abs. of Std.}}$$

Abs=absorbance

Conc.=concentration

Std=standard.

DETERMINATION OF TRIGLYCERIDE (enzymatic colorimetric method) of Trinder [25].

Assay procedure;

1000µl of triglyceride liquor and 10µl of sample, standard and control were added into test tubes respectively, and incubated for 10 minutes at 37°C. The absorbance of standard and test were read (measured) against reagent blank within 60 minutes at 500nm wavelength.

$$\text{Conc. of triglyceride} = \frac{\text{Abs. of test} \times \text{Conc. of Std.}}{\text{Abs. of Std.}}$$

Abs=absorbance

Conc.=concentration

Std=standard.

DETERMINATION OF HIGH-DENSITY LIPOPROTEIN (HDL)

ASSAY PROCEDURE; 100µl of precipitating reagent and 100µl of sample were added into dry test-tube and incubated at room temperature for 5mins. The solution was centrifuged at 3000 rpm (revolution per minute). 50µl of the sample supernatant, standard solution and distilled water (as blank solution) were added into 1000µl of cholesterol reagent respectively. The solutions were incubated for 15mins at 25°C. The absorbance was read at 505nm against the blank within 60mins.

$$\text{Concentration of HDL cholesterol (mg/dl)} = \frac{\text{Abs. T} \times \text{Conc. Std} \times 2}{\text{Abs. Std}}$$

CALCULATION OF LDL (Freidewald's Formula)

$$\text{Total cholesterol} - \frac{(\text{triglyceride})}{5} - \text{HDL Cholesterol}$$

RESULTS

Table 1: Mean ± SD values of lipid profile of the studied population.

Parameters	Diabetic patients (n=100)	Controls (n=50)	t-value	p-value
Cholesterol (mmol/l)	4.96±0.91	4.48±0.41	3.584	0.001
Lower 95% C.I	0.21	0.27		
Upper 95% C.I	0.75	0.69		
Triglyceride (mmol/l)	1.84±0.42	1.63±0.09	3.420	0.001
Lower 95% C.I	0.08	0.12		
Upper 95% C.I	0.33	0.29		
High density Lipoprotein (mmol/l)	1.63±0.43	1.99±0.32	-5.204	0.001
Lower 95% C.I	-0.49	-0.48		
Upper 95% C.I	-0.22	-0.23		
Low density	2.40±0.66	1.67±0.22	7.574	0.001

Lipoprotein(mmol/l)				
Lower 95% C.I	0.54	0.58		
Upper 95% C.I	0.92	0.88		
Atherogenic index	0.053±0.01	-0.087±0.72		0.001

Values of serum lipid profile levels of the studied population

The serum Cholesterol, Triglyceride and low-density lipoprotein were statistically higher ($p < 0.001$, $p = 0.001$, $p < 0.001$ respectively) in the studied population compared with controls. The serum High density lipoprotein was statistically lower ($p < 0.001$) in the studied population compared with control. Atherogenic index was statistically higher ($p = 0.001$) in the studied population compared with control.

DISCUSSION

In this present work, the serum Cholesterol, Triglyceride, Low density lipoprotein, blood platelet and prothrombin time levels were higher in studied population compared with controls. There was higher level of hemoglobin in male studied population compared with female studied population. The higher levels of Cholesterol, Triglyceride and Low-density lipoprotein, could possibly be due to altered metabolism of triglyceride-rich lipoproteins which is crucial in the pathophysiology of the atherogenic dyslipidemia of diabetes. Alterations include both increased hepatic secretion of Very Low-Density Lipoprotein (VLDL) and impaired clearance of VLDL and intestinally derived chylomicrons. An important consequence of retarded clearance is prolonged plasma retention of both VLDL and postprandial chylomicrons as partially lipolyzed remnant particles.

CONCLUSION

From the study since cholesterol, triglyceride, low density lipoprotein was higher in diabetic patients and are associated with risk factors as coronary artery disease, cardiovascular disease, atherosclerotic diseases, atherothrombotic complications and deficiency of vitamin k, care has to be taken in managing the disease.

REFERENCES

- Janez A, Guja C, Mitakou A. Insulin Therapy in Adults with Type 1 Diabetes Mellitus: a Narrative Review. *Diabetes Therapy*, 2020; 11: 387-409.
- Ifediora AC, Obeagu EI, Akahara IC, Eguzouwa UP. Prevalence of urinary tract infection in diabetic patients attending Umuahia health care facilities. *J Bio Innov*, 2016; 5(1): 68-82. [links/5ae45fdaca272ba507eb3c3/PREVALENCE-OF-URINARY-TRACT-INFECTION-IN-DIABETIC-PATIENTS-ATTENDING-UMUAHIA-HEALTH-CARE-FACILITIES.pdf](https://doi.org/10.5966/jbioinnov.2016.0501006).
- Ugwu OP, Alum EU, Okon MB, Aja PM, Obeagu EI, Onyeneke EC. Ethanol root extract and fractions of *Sphenocentrum jollyanum* abrogate hyperglycaemia and low body weight in streptozotocin-induced diabetic Wistar albino rats. *RPS Pharmacy and Pharmacology Reports*, 2023; 2(2): rqad010.
- Obeagu EI, Obeagu GU. Utilization of Antioxidants in the management of diabetes mellitus patients. *J Diabetes Clin Prac*, 2018; 1(102): 2. [links/5b6c2dec92851ca65053b74e/Utilization-of-Antioxidants-in-the-Management-of-Diabetes-Mellitus.pdf](https://doi.org/10.5966/jdiabetes.2018.0102002).
- Obeagu EI, Okoroiwu IL, Obeagu GU. Some haematological variables in insulin dependent diabetes mellitus patients in Imo state Nigeria. *Int. J. Curr. Res. Chem. Pharm. Sci.*, 2016; 3(4): 110-7. [links/5ae4abee458515760ac07a13/Some-haematological-variables-in-insulin-dependent-diabetes-mellitus-patients-in-Imo-state-Nigeria.pdf](https://doi.org/10.5966/ijcrps.2016.0304011).
- Nwakuilite A, Nwanjo HU, Nwosu DC, Obeagu EI. Evaluation of some trace elements in streptozocin induced diabetic rats treated with Moringa oleifera leaf powder. *WJPMR*, 2020; 6(12): 15-8. [links/5fcb587092851c00f8516430/EVALUATION-OF-SOME-TRACE-ELEMENTS-IN-STREPTOZOCIN-INDUCED-DIABETIC-RATS-TREATED-WITH-MORINGA-OLEIFERA-LEAF-POWDER.pdf](https://doi.org/10.5966/wjpmr.2020.0612015).
- Desai S, Deshmukh A. Mapping of Type 1 Diabetes Mellitus. *Current Diabetes Review*, 2020; 16: 438-41.
- Anyiam AF, Obeagu EI, Obi E, Omosigho PO, Ironi EA, Arinze-Anyiam OC, Asiyah MK. ABO blood groups and gestational diabetes among pregnant women attending University of Ilorin Teaching Hospital, Kwara State, Nigeria. *International Journal of Research and Reports in Hematology*, Jun. 21, 2022; 5(2): 113-21.
- Okafor CJ, Yusuf SA, Mahmoud SA, Salum SS, Vargas SC, Mathew AE, Obeagu EI, Shaib HK, Iddi HA, Moh'd MS, Abdulrahman WS. Effect of Gender and Risk Factors in Complications of Type 2 Diabetic Mellitus among Patients Attending Diabetic Clinic in Mnazi Mmoja Hospital, Zanzibar. *Journal of Pharmaceutical Research International*, May. 25, 2021; 33(29B): 67-78.
- Galano ES, Yusuf SA, Ogbonnia SO, Ogundahunsi OA, Obeagu EI, Chukwuani U, Okafor CJ, Obianagha NF. Effect of Extracts of Kigelia Africana Fruit and Sorghum Bicolor Stalk on the Biochemical Parameters of Alloxan-Induced Diabetic Rats. *Journal of Pharmaceutical Research International*, Apr. 26, 2021; 33(25B): 86-97.
- Kama SC, Obeagu EI, Alo MN, Ochei KC, Ezugwu UM, Odo M, Ikpeme M, Ukeekwe CO, Amaeze AA. Incidence of Urinary Tract Infection among Diabetic Patients in Abakaliki Metropolis. *Journal of Pharmaceutical Research International*, Nov. 17, 2020; 32(28): 117-21.

12. Nwakulite A, Obeagu EI, Eze R, Vincent CC, Chukwurah EF, Okafor CJ, Ibekwe AM, Adike CN, Chukwuani U, Ifionu BI. Evaluation of Catalase and Manganese in Type 2 Diabetic Patients in University of Port Harcourt Teaching Hospital. *Journal of Pharmaceutical Research International*, Jun. 3, 2021; 40-5.
13. Cieluch A, Uruska A, Zozulinska-Ziolkiewicz D. Can We Prevent Mitochondrial Dysfunction and Diabetic Cardiomyopathy in Type 1 Diabetes Mellitus? Pathophysiology and Treatment Options. *International Journal of molecular Science*, 2020; 21: 2852.
14. Nwakulite A, Obeagu EI, Nwanjo HU, Nwosu DC, Nnatuanya IN, Vincent CC, Amaechi CO, Ochiabu O, Barbara MT, Ibekwe AM, Okafor CJ. Studies on Pancreatic Gene Expression in Diabetic Rats Treated with Moringa oleifera Leaf. *Journal of Pharmaceutical Research International*, May. 4, 2021; 33(28A): 78-86.
15. Nwosu DC, Nwanjo HU, Obeagu EI, Ugwu GU, Ofor IB, Okeke A, Ochei KC, Kanu SN, Okpara KE. EVALUATION OF LIPOPROTEIN A AND LIPID TETRAD INDEX PATTERN IN DIABETIC PATIENTS ATTENDING METABOLIC CLINIC IN THE FEDERAL MEDICAL CENTRE, OWERRI, IMO STATE. **World Journal of Pharmacy and Pharmaceutical Sciences**, 2015; 4(3): 126-140
[links/5a4fd1a00f7e9bbc10526b54/EVALUATION-OF-LIPOPROTEIN-A-AND-LIPID-TETRAD-INDEX-PATTERN-IN-DIABETIC-PATIENTS-ATTENDING-METABOLIC-CLINIC-IN-THE-FEDERAL-MEDICAL-CENTRE-OWERRI-IMO-STATE.pdf](https://www.ejbps.com/links/5a4fd1a00f7e9bbc10526b54/EVALUATION-OF-LIPOPROTEIN-A-AND-LIPID-TETRAD-INDEX-PATTERN-IN-DIABETIC-PATIENTS-ATTENDING-METABOLIC-CLINIC-IN-THE-FEDERAL-MEDICAL-CENTRE-OWERRI-IMO-STATE.pdf).
16. Ezema GO, Omeh NY, Egbachukwu S, Agbo EC, Ikeyi AP, Obeagu EI. Evaluation of Biochemical Parameters of Patients with Type 2 Diabetes Mellitus Based on Age and Gender in Umuahia. *Asian Journal of Dental and Health Sciences*, Jun. 15, 2023; 3(2): 32-6.
<http://ajdhs.com/index.php/journal/article/view/43>.
17. Adu ME, Chukwuani U, Ezeoru V, Okafor CJ, Amaechi CO, Vincent CC, Obeagu GU, Eze R, Nnatuanya IN, Nwosu DC, Nwanjo HU. Studies on molecular docking of moringa oleifera leaf phytochemical constituents on alpha glucosidase, alpha amylase and dipeptidyl peptidase. *Journal of Pharmaceutical Research International*, May. 7, 2021; 33(28A): 239-45.
18. Ezugwu UM, Onyenekwe CC, Ukibe NR, Ahaneku JE, Obeagu EI. Plasma Level of Macromolecules and Mathematical Calculation of Potential Energy in Type 2 Diabetic Individuals at NAUTH, Nnewi, Nigeria. *Journal of Pharmaceutical Research International*, Nov. 2, 2021; 33(47B): 242-8.
19. Ripsin CM, Kang H, Urban RJ. Management of blood glucose in type 2 diabetes mellitus (PDF). *American Family Physician*, 2013; 79(1): 29-36.
20. Nwakulite A, Obeagu EI, Eze R, Ugochi VE, Vincent CC, Okafor CJ, Chukwurah EF, Unaeze BC, Amaechi CO, Okwuanaso CB, Chukwuani U. Estimation of Serum Glutathione Peroxidase in Streptozotocin Induced Diabetic Rat Treated with Bitter Leaf Extract. *Journal of Pharmaceutical Research International*, Jun. 7, 2021; 33(30B): 200-6.
21. Okoroiwu IL, Obeagu EI, San Miguel HG, Bote SA, Obeagu GU. Characterisation of HLA-DR antigen in patients type 1 diabetes mellitus in patient attending a tertiary hospital in Enugu, south-east Nigeria. *ACADEMIC JOURNAL*, 2023.
22. Okoroiwu IL, Obeagu EI, Obeagu GU, Chikezie CC, Ezema GO. The prevalence of selected autoimmune diseases. *Int. J. Adv. Multidiscip. Res.*, 2016; 3(3): 9-14.
23. Nwakulite A, Nwanjo HU, Nwosu DC, Obeagu EI. EVALUATION OF ENZYME ANTIOXIDANTS IN STREPTOZOCIN INDUCED DIABETIC RATS TREATED WITH MORINGA OLEIFERA LEAF POWDER. *European Journal of Biomedical*, 2020; 7(11): 285-8.
24. Nwosu DC, Nwanjo HU, Opara AU, Ofor IB, Obeagu EI, Ugwu GU, Ojiegbe GC, Nnorom RM, Nwokike GI, Okpara KE, Ochei KC. EVALUATION OF C-REACTIVE PROTEIN, SELENIUM AND GLYCOSYLATED HAEMOGLOBIN LEVELS IN DIABETIC PATIENTS ATTENDING METABOLIC CLINIC IN THE FEDERAL MEDICAL CENTRE, OWERRI, IMO STATE. **World Journal of Pharmacy and Pharmaceutical Sciences**, 2015; 4(3): 141-152.
https://www.academia.edu/download/38320132/NWOSU_EMMA_9.pdf.
25. Trinder P. Enzymatic Colorimetric determination of triglyceride by GOP-PAP method, 1969.